

**Nigerian Journal of
Animal Science**
(Formerly Tropical Journal of Animal Science)

22



Volume 21 Number 3,
2019.

Published by the
Animal Science Association of Nigeria

ISSN 1119-4308

Impact of black plum (*Vitex doniana*) leaf meal on blood biochemistry, hormone and cholesterol level of West African dwarf goat-bucks

Okukpe K. M*, Lawal M.O., Sanni K.M., Adeyina A.O., Alli O.I., DeCampos J.S. and Aderibigbe T. A.

Reproductive Physiology Unit, Department of Animal Production, Faculty of Agriculture.

*Corresponding author: okukpekehinde@yahoo.com; Phone number: +2348066716145

Target Audience: Animal scientists, Extension officers, Goat producers, Meat processors

Abstract

Fifteen (15) West African Dwarf (WAD) goat-bucks with average initial weights of 10 ± 0.53 kg were used in an experiment to investigate the effect of varying levels of *Vitex doniana* leaf meal on the blood profile, hormone and cholesterol level of WAD goat-bucks. The animals were randomly assigned to five dietary treatments consisting of graded levels of *Vitex doniana* in a Completely Randomized Design experiment which lasted twenty-eight (28) days. At the end of the feeding trial, blood samples were collected from the jugular veins of the animal for the determination of haematological and serum biochemical characteristics using standard laboratory methods. The *Vitex doniana* leaf had a crude protein of 8.53, crude fibre of 7.13, crude fat of 6.44 and ash of 8.26 percent and were moderately high in flavonoids and anthraquinones-saponin 6.48mg/100g with slight presence of alkaloids, tannins, terpenoids, steroids and anthraquinones. The effects of graded levels of *Vitex doniana* leaf meal on haematology of WAD goat-bucks reveals that red blood cell (RBC) significantly ($p < 0.05$) increase with concurrent increase in *Vitex doniana* leaf meal, whereas the haemoglobin (Hb) and packed cell volume (PCV) reduced with the addition of the leaf meal. Other parameters were not significantly ($p > 0.05$) different from the control, although it varies with increase in the leaf meal addition. The serum biochemistry of West African dwarf bucks showed that total protein (TP), albumin (ALB), urea and glucose significantly increased with increase in *Vitex doniana* leaf meal. Alkaline phosphatase (ALP), Alanine aminotransferase (ALT) and Aspartate aminotransferase (AST) was significantly ($p < 0.05$) influenced by the increase in *Vitex doniana* leaf meal inclusion. The ALT and AST were not significantly ($p > 0.05$) different from the control in treatment D. Phytochemical results showed *Vitex doniana* has high content of phenol, flavonoid and saponin; it significantly increase total cholesterol and low density lipoprotein with a reduction in high density lipoprotein with 20g/kg feed while triglyceride and high density lipoprotein were high with 10g/kg feed. It significantly reduced luteinizing hormone while causing an increase in follicle stimulating hormone, testosterone and prolactin with 15g/kg. The administration of 15g inclusion level of *Vitex doniana* leaf meal increases the reproductive performance while administration of 10g inclusion level helped to lower the cholesterol level in WAD bucks with no conspicuous adverse effects and their health status was not compromised. All the haematological and serum biochemical characteristics of the bucks were within the normal/ standard blood ranges for apparently healthy bucks. The study therefore concludes that farmers can supplement feed with up to 20% *Vitex doniana* leaf meal in formulated diets meant for goats, without fear of compromising haematopoietic processes.

Key words: Blood, Feed, leaf-meal, WAD goat-bucks.

Description of Problem

Production of ruminant animals in developing African countries is often

characterized with low level of efficiency. The apparent inefficiency of ruminant production has been attributed to unfavourable climate

disease prevalence and feed shortage especially during dry seasons. The first two factors tend to encourage adoption of intensive ruminant farming system which can further aggravate the problem of inadequate feed supply. The cost of feed accounts for about 60% of total intensive production cost compared to 40% value under extensive production system (2). A large reduction in feed cost is achievable by the use of unconventional feed resources such as fodder

used in profiles, e dietary experiment d from the characteristics are of 7.15 nols 96.14 raquinones, eals that red- ra leaf meal, he leaf meal, t varies with wed that total x doniana leaf nsferase (AST), The ALT and al results shows otal cholesterol hile triglyceride hormone while th 15g/kg. The ve performance,) bucks with no natological, and ood ranges for with up to 26% haematopoietic

efficiency. The ant. production urable climate.

Table 1: Composition of experimental basal diet

Ingredients	% Inclusion
Maize	30.00
Groundnut cake	14.00
Wheat offal	33.00
Rice bran	10.00
Palm kernel cake	10.00
Table Salt	1.00
Premix	0.25
Limestone	1.00
Bone meal	0.75
Total	100.00

Materials and Methods

Site and Period of the Study

Feeding trials and laboratory analyses were conducted in the Livestock unit of the University Farm and the Animal Production Laboratory respectively at the Department of Animal Production, University of Ilorin located on Latitude 08° 29'N; Longitude 004° 35' E; Elevation 308 m, having a mean annual rainfall of 98.75 mm that is spread largely over the months of May to October.

Collection and Processing of *Vitex doniana* Leaves

Fresh leaves were collected from *Vitex doniana* (Black plum) trees on the University of Ilorin farm during the dry season months of November and December. Leaf samples were identified at the Department of Plant Biology Herbarium, Faculty of Life Science, University of Ilorin and a sample deposited with assigned voucher number. The leaves were subsequently air-dried for 10 days and kept inside jute bags in a well ventilated open shed that was protected from rain and direct rays of sunlight. The leaves were later processed by grinding in a hammer mill to pass through 2 mm sieve and kept in jute bags for subsequent feed formulation. Samples of the leaves were analyzed for gross energy, dry matter and chemical components (Table 2).

The aim of the present work seeks to explore the potential of *Vitex doniana* (black plum) leaves, which presently are of little economic value for goat feeding under intensive system of production.

Table 2: Proximate composition of experimental diets and *Vitex doniana* leaf meal

Parameter (%)	A	B	C	D	E	V
Dry matter	84.62	91.46	90.88	90.98	91.51	92.75
Moisture content	15.38	8.54	9.12	9.02	8.49	7.25
Crude fat	3.94	4.98	2.91	4.39	5.42	6.44
Crude protein	16.19	16.59	19.69	16.79	15.31	8.53
Crude fibre	6.05	13.40	8.35	9.03	9.14	7.15
Ash content	8.69	10.23	8.33	9.50	8.56	8.26

Where A is the control (basal diet without *Vitex doniana* leaf meal), B (basal +5g *V. doniana*), C (basal +10g *V. doniana*), D (basal +15g *V. doniana*), E (basal +20g *V. doniana*), V (*Vitex doniana* leaf meal)

Feed Formulation and Processing

The leaves were mixed with crushed maize, soy bean meal, bone meal and sodium chloride in the different inclusion levels of 0, 5, 10, 15 and 20g/kg of leaves to feed as indicated in Table 1 and 2. The resultant feeds were processed into crumbles to reduce dustiness, facilitate handling and encourage intake by animals. Each feed was wetted with about 2% (v/w) clean drinking water and sundried on concrete slab for 5 – 6 hours.

Experimental Design and Animal Management

A 28 – day growth and nutrient digestibility trial was conducted with fifteen weanling male West African Dwarf goats of average initial live weight of 10.50 ± 0.34 kg. The goats were divided into five dietary treatments of three bucks each in a completely randomized design. Animals in treatment A (control) were fed formulated diet without *Vitex doniana* leaf meal while the other four treatment (B, C, D and E) had *Vitex doniana*

leaf meal in varying inclusion level of 5, 10, 15 and 20g/kg of feed respectively. Each animal were fed 300g/day of the formulated feed throughout the experimental period. *Panicum maximum* were also provided in abundance for the bucks.

On the 28th day, blood was collected by jugular vein puncture with hypodermic needles and syringe from three animals per treatment into two (2) clean sample bottles, one with anticoagulant, Ethylene Diamine tetra Acetate (EDTA) for haematological analysis and the other without EDTA for serum biochemical analysis. All haematological parameters were determined by conventional laboratory methods of (3). Serum biochemical parameters were assayed using standard laboratory methods (12). All data were analyzed using the analysis of variance (ANOVA) procedure following a completely randomized model (16) and the levels of significance were determined using the Duncan's Multiple Range Test.

Table 3: Phytochemical composition of *Vitex doniana* leave meal

PHYTOCHEMICALS	QUALITATIVE	QUANTITATIVE (mg/100g)
Alkaloids	+	2.61
Saponins	++	6.48
Tannins	+	1.45
Flavonoids	++	20.82
Terpenoids	+	0.21
Phenols	+++	96.14
Steroids	+	2.02
Anthraquinones	+	0.04

KEY: + PRESENT; ++ MODERATELY PRESENT; +++ HIGHLY PRESENT

Results a
Vitex don
high in
necessary
against n
diets of
caused a
blood ce
Vitex d
haemogl
(PCV) and
addition
paramete
significa

Table 4:
dwarf b
Paramete

Red blood
cells $\times 10^{12}$
Haemoglobin
(g/dl)
Packed ce
Volume (%)
White blo
cells $\times 10^9$
Neutroph
Lymphoc
Monocyt
Eosinoph

a, b, c, d
V. doniana
V. donia

Red-b
obtain
V. donia
signifi
treatm
in tre
treatm
(1) in
reporte
could

Results and discussion

Vitex doniana leaf meal was observed to be high in flavonoids and phenols which are necessary immunostimulants for defence against microbial invasion (13). Its inclusion in diets of West African Dwarf goat-bucks caused a significant increase ($p < 0.05$) in red blood cells (RBC) with concurrent increase in *Vitex doniana* leaf meal. However, the haemoglobin (Hb) and packed cell volume (PCV) significantly reduce with increased addition of the leaf meal (Table 4). Other parameters though similar to the control was significantly different ($p < 0.05$) from the

treatment D, although it varies with increase in the leaf meal addition. Table 5 shows the effects of graded levels of *Vitex doniana* leaf meal on serum biochemistry of WAD goat-bucks. The total protein (TP), albumin (ALB), urea and glucose significantly increased with increase in *Vitex doniana* leaf meal. Alkaline phosphatase (ALP), Alanine aminotransferase (ALT) and Aspartate aminotransferase (AST) was significantly influenced by the increase in *Vitex doniana* leaf meal inclusion. The ALT and AST were not significantly different ($p > 0.05$) from the control in treatment D.

Table 4: Effects of graded levels of *Vitex doniana* leaf meal on haematology of West African dwarf bucks

Parameters	A	B	C	D	E	±SEM
Red blood cells $\times 10^{12}/L$	12.68 ^d	12.88 ^c	13.18 ^b	13.22 ^b	13.89 ^a	0.03
Haemoglobin (g/dl)	9.86 ^b	10.70 ^a	10.63 ^a	9.22 ^c	9.36 ^c	0.26
Packed cell volume (%)	26.00 ^b	28.33 ^a	29.00 ^a	24.00 ^c	24.33 ^c	1.32
White blood cells $\times 10^9/L$	13.33 ^a	13.00 ^{ab}	13.00 ^{ab}	12.56 ^b	12.78 ^{ab}	1.81
Neutrophils (%)	21.67 ^{ab}	20.00 ^b	23.33 ^a	23.33 ^a	23.33 ^a	1.84
Lymphocyte (%)	83.70 ^{ab}	90.03 ^a	78.45 ^b	77.04 ^b	74.07 ^b	1.23
Monocytes (%)	2.33 ^b	3.00 ^a	1.33 ^c	1.33 ^c	1.33 ^c	0.37
Eosinophil (%)	0.00 ^b	1.00 ^a	0.00 ^b	0.00 ^b	0.00 ^b	0.87

a, b, c, d – means along the rows with different superscripts differ significantly ($p < 0.05$). A (basal diet without *Vitex doniana* leaf meal), B (basal + 5g *V. doniana*), C (basal + 10g *V. doniana*), D (basal + 15g *V. doniana*), E (basal + 20g *V. doniana*).

Red blood cell (RBC) of $13.89 \pm 0.03 \times 10^{12}/L$ obtained for bucks that received 20g *Vitex doniana* leaf meal in treatment E was significantly ($p < 0.05$) higher than all other treatments. RBC values of $12.88 \pm 0.03 \times 10^{12}/L$ in treatment A and $13.89 \pm 0.03 \times 10^{12}/L$ in treatment E were in contrast with the report of (1) in Saanen goats but were within the range reported by (7) for West African Dwarf goat. It could be said that RBC value changes with

breed. It therefore shows that *Vitex doniana* has no negative effect on the health status of the animal. Haemoglobin concentration (g/dl) and PCV (%) were significantly higher ($p < 0.05$) in treatment B and C but lower in treatment C and D. The haemoglobin concentration and PCV were within the normal range reported by (7) and (17) for goats respectively. Increase in haemoglobin implies that the animals were able to transport oxygen

to tissues for oxidation of ingested food so as to release energy for the other body functions (14). The WBC counts were in contrast with the report of (1) in Saanen goat. This difference could be as a result of breed, environmental factor and age of the animal. Neutrophil, the most abundant of white blood cell type increases from treatment B to C but become constant with additional increase in *Vitex doniana* leaf meal. Lymphocytes are the largest type of white blood cell and it is a-

granulocyte. It was significant ($p < 0.05$) reduced with increase in the *Vitex doniana* leaf meal in the diet. Eosinophil was only significant in treatment B. Eosinophils are responsible for fighting infections of parasitic worms. These cells release toxins that kill the worms and are also involved in the inflammatory response when there is an allergic reaction. The value was within the normal range reported by (7) and (20).

Total (P<0.05) with the c improve buffer in t and a car such as minerals. calcium, the blood l Albumin v protein wa the higher This impli 20g (impr creatinine metabolism respectively body-like shows a r significant was within Hence, the on bucks. at treatme and D. W range, it while hi hyperglyc (ALP) inc leaf me aminotran range rep was an inc (AST) le inclusion range (12 and organ elevation would hav The treatment with treat be due compensa

Table 5: Effects of graded levels of *Vitex doniana* leaf meal on serum biochemistry of West African dwarf bucks

Parameters	A	B	C	D	E	±SEM
Total protein (g/L)	10.00 ^b	11.67 ^a	9.65 ^{ab}	9.15 ^b	11.33 ^a	1.19
Albumin (g/L)	3.67 ^c	3.77 ^b	3.67 ^c	3.00 ^d	4.44 ^a	0.35
Urea (mmol/L)	2.00 ^d	2.53 ^c	3.10 ^b	2.07 ^d	5.73 ^a	0.26
Creatinine (μmol/L)	0.24 ^d	0.39 ^a	0.24 ^d	0.27 ^c	0.31 ^b	1.04
Glucose (mmol/L)	0.57 ^b	0.87 ^a	0.33 ^c	0.30 ^c	0.83 ^a	0.05
ALP (IU/L)	86.00 ^c	68.67 ^d	98.00 ^b	108.33 ^a	108.33 ^a	2.83
ALT (IU/L)	27.67 ^a	26.00 ^b	21.67 ^c	28.00 ^a	17.67 ^d	1.19
Total cholesterol (mmol/L)	0.65 ^{ab}	0.45 ^b	0.70 ^{ab}	0.65 ^{ab}	1.20 ^a	0.14
Triglyceride (mmol/L)	0.10 ^b	0.15 ^b	0.45 ^a	0.10 ^b	0.15 ^b	0.02
Low-density lipoprotein (mmol/L)	0.30 ^b	0.30 ^b	0.45 ^{ab}	0.40 ^b	0.70 ^a	0.06
Luteinizing hormone (MIU/ml)	0.40 ^a	0.25 ^b	0.15 ^{bc}	0.10 ^c	0.10 ^c	0.03
Follicle-stimulating hormone (MIU/ml)	0.50 ^b	1.50 ^b	3.25 ^a	4.75 ^a	4.00 ^a	0.39
Testosterone (ng/dl)	5.50 ^b	2.20 ^c	0.30 ^d	7.10 ^a	0.40 ^d	0.34
Prolactin (μg/L)	0.50 ^b	0.25 ^b	0.50 ^b	0.90 ^a	0.60 ^{ab}	0.19

a, b, c, d – means along the rows with different superscripts differ significantly ($p < 0.05$): A (basal diet without *Linum donianum* leaf meal), B (basal + 5g l% *doniana*), C (basal + 10g l% *doniana*), D (basal + 15g l% *doniana*), E (basal + 20g l% *doniana*), ALT- Alanine aminotransferase, ALP- Alkaline phosphatase and AST- Aspartate aminotransferase.

ry of West

±SEM

1.19

0.35

0.26

1.04

0.05

2.88

1.19

0.14

0.02

0.06

0.03

0.89

0.34

0.19

without Vitex
E (basal) = 20g
ansferase

Total protein value was significantly high ($p < 0.05$) in most of the treatments compared with the control. *Vitex doniana* leaf seems to improve the total protein which serves as buffer in the maintenance of Acid-base balance and a carrier of essential blood constituents such as hormone, vitamins and certain minerals. Plasma protein helps to transport calcium, phosphorus and other substances in the blood by attachment to the albumin (6; 13). Albumin which is most abundant serum total protein was highest in treatment E which had the highest inclusion of *Vernonia doniana*. This implies that *Vitex doniana* leaf meal at 20g improve Albumin level. Urea and creatinine are waste products of protein metabolism and waste generated from creatine respectively. High level of urea is toxic to the body likewise the high value of creatinine shows a renal dysfunction. Though, urea was significantly high in treatment E and C, but was within the normal range reported by (7). Hence, the leaf meal has no deleterious effect on bucks. Glucose level was significantly high at treatments B and E but low in treatment C and D. When glucose is lower than the normal range, it is an indication of hypoglycaemia while higher levels are indication of hyperglycaemia (13). Alkaline phosphatase (ALP) increased with increase in level of the leaf meal inclusion while Alanine aminotransferase (ALT) was within the normal range reported by (1) in Saanen goats. There was an increase in Aspartate aminotransferase (AST) level with increase in leaf meal inclusion but it was still within the normal range (12). These enzymes are tests of liver and organ functions of the animal and any elevation above or below the normal range would have indicated organ dysfunction.

The total cholesterol was highest in treatment E but low in the other treatments with treatment B having the lowest. This may be due to the fact that the body also compensates for any absorption of additional

cholesterol by reducing cholesterol synthesis as reported by (10). For these reasons, seven to ten hours after ingestion of cholesterol, blood levels will show little if any effect on total body cholesterol content or concentrations of cholesterol in the blood. However, during the first seven hours after ingestion of cholesterol, the levels significantly increase (13). The observed increase in total cholesterol, triglycerides and low-density lipoprotein could be attributed to its health and medicinal effects of three groups of active phytochemicals (tannins, saponins, alkaloids). Saponin (in excess) causes hypocholesterolaemia because it binds cholesterol making it unavailable for absorption (15).

There were significant differences ($p < 0.05$) in luteinizing hormone (LH) value and these ranges from 0.10 ± 0.03 in treatment D and E to 0.40 ± 0.03 in the control. LH has been reported to stimulate testosterone output in a variety of *in-vitro* systems including perfused testes, incubations of testicular slices, minces or homogenates, incubations of entire decapsulated testes and in suspensions of purified Leydig cells (5).

Follicle stimulating hormones (FSH) significantly increase with increased level of *Vitex doniana* leaf meal. This support the work of (8) that increase in serum estradiol levels cause a decrease in FSH production by inhibiting gonadotrophin releasing hormone (GnRH) production in the hypothalamus. A decrease in serum FSH level would cause the smaller follicles in the ovary to undergo atresia as they lack sufficient sensitivity to FSH to survive (18). The significantly high levels of testosterone in treatment D (7.10 ± 0.34 ng/dl) could be as a result of increased *Vitex doniana* leaf meal on FSH/LH secreting organ. Testosterone is essential for muscle lay-down, health and well-being as well as the prevention of osteoporosis (19). Prolactin level was significantly high with increase in *Vitex*

doniana leaf meal with the highest in treatment D (15g inclusion level). Prolactin has been reported to enhance spermatogenesis by augmenting the effect of endogenous or exogenous LH on testicular steroidogenesis rather than potentiating the effect of androgens on the seminiferous epithelium (4).

Conclusion and Application

In conclusion,

1. *Vitex doniana* leaf meal has a valuable non-conventional feedstuff potential; Livestock farmers can use it as a feed supplement in the diet of West African Dwarf bucks with up to 20g/kg concentrate feed, without fear of compromising haematopoietic processes and blood chemistry.
2. Optimum performance in terms of moderating cholesterol and reproductive function can be achieved with a 15g inclusion of *Vitex doniana* leaf meal in concentrate diets.
3. Further studies should be done on the mechanism for effecting the changes on the animal and the residual effect on the animal product.

References

1. Afyon Kocatepe. (2012). Reference values for haematological and biochemical parameters in Saanen goats breeding in Afyon karahisar Province, *Kocatepe Veterinary Journal*, 5(1): 7-11.
2. Atteh, J. O. (2002). Principles and Practices of Livestock Feed and Manufacturing, Ilorin. Adlek printers, Ilorin, Nigeria.
3. Baker, F. S. and Silverton, R. E. (1982). Introduction to medical laboratory technology, 5th ed. Butterworth S. C. London Pp481-494.
4. Bartke A., Hafiez A.A., Bex F.J. and Dalterio S. (1978). Hormonal Interactions in regulation of androgen secretion. *Biology of Reproduction*, 18: 44-54.
5. Catt K.J. and Dafau M.L. (1976). Basic concepts of the mechanism of action of peptide hormones. *Biology of Reproduction*, 14: 1-15.
6. Dacie, J. V. and Lewis, S. M. (2001). Practical Haematology. 9th Edition, 362. Churchill Livingstone, New York, page 633.
7. Daramola, J. O., Adeloye, A. A., Fatoba T. A. and Soladoye, A. O. (2005). Haematological and biochemical parameters of West African Dwarf goats. *Livestock Research for Rural Development*, 17 (8): 1-8.
8. El Safoury S and Bartke A. (1974). Effects of follicle-stimulating hormone and luteinizing hormone on plasma testosterone levels in hypophysectomized and in-tact immature and adult male rats. *Journal of Endocrinology*, 61: 193-198.
9. Frances, I. O., Thomas, M. O., Gabrel, O. (2013). Effect of processing methods on the chemical composition of *Vitex doniana* leaf and leaf products. *Food Science & Nutrition*, 1(3): 241-245.
10. Lecerf JM, de Lorgeril M (2011). "Dietary cholesterol: from physiology to Cardiovascular risk". *British Journal of Nutrition*, 106 (1): 6-14.
11. Makun, H., Otaru, S.M., Dung, D. (2013). Effect of management practices on milk yield and live weight changes in indigenous breeds of goats supplemented with groundnut haulms and concentrate in sub humid zone of Nigeria. *Sokoto Journal of Veterinary Science*, 11(1): 45-50.
12. Ochei, J. and Kolhatkar, A. (2007). *Medical Laboratory Sciences Theory and Practice*, 6th edn. McGraw Hill Pub. Co. Ltd., New York, U. S. A., 1338PP.
13. Okukpe, K. M., Adeyina, O. A., Olatunde, O. A. and Alli O. I. (2015). Fundamentals of Anatomy and

- Physiology of Farm Animals. First Edition, Olad publication, Nigeria, page 59-87.
4. Olorunnisomo, O. A., Ewuola, E. O. and Lawal, T. T. (2012). Intake and blood metabolites in Red Sokoto Goats fed Elephant Grass and Cassava Peel Silage. *Journal of Animal Production Advances*, 2(9): 420-428.
 5. Soetan K.O. and Oyewole O.E. (2009). The need for adequate processing to reduce the anti-nutritional factors in animal feeds, a review. *African Journal of Food Science*, 3(9), 223-232.
 6. Steel, R. G. D and Torrie, J. H. (1980). *Principles and procedures of statistics. A Biometrical Approach*, 2nd Edition, McGraw Hill Book Co. Inc. New York. USA. 481Pp.
 7. Tambuwal, F. M., Agale B. M. and Bangana, A. (2002). Haematological values of apparently healthy Red Sokoto Goats. *Proceeding of 27th Annual Conference of the Nigerian Society for Animal Production (NSAP)*, March 2002. Federal University of Technology, Akure, Nigeria. Page 50-53.
 18. Thorner, M. O., Edwards C. R. W., Harker, J. P., Abraham, G. and Besser, G. M. (1977). Prolactin and Gonadotropin interaction in the male. In: *The testis in normal and infertile men*. (P. Troen and H. R. Nakin. eds), Raven Press, New York, page 351-366.
 19. Tuck SP, Francis RM (2009). "Testosterone, bone and osteoporosis". *Frontiers of Hormone Research*, 37: 123-32.
 20. Zamiri, M. J. and Heidari, A. H. (2006). Reproductive characteristics of Ragini male goats of Kerman province in Iran. *Animal Reproductive Science*, 16: 540-561.