



**VOLUME 2, 2019**  
**ISSN: 2695 - 1959**



# **OSUN GEOGRAPHICAL REVIEW**

**JOURNAL OF THE DEPARTMENT OF GEOGRAPHY,**  
**OSUN STATE UNIVERSITY, STATE OF OSUN, NIGERIA**

**Published by the**  
Department of Geography, Osun State University,  
State of Osun, Nigeria



Hand Copy  
Law Kgun O  
Department of Geography  
University of Ilorin

---

# OSUN GEOGRAPHICAL REVIEW

Journal of the Department of Geography,  
Osun State University, State of Osun, Nigeria

---

Volume 2, 2019

ISSN: 2695 - 1959

**Editor-in-Chief:**  
Prof. Olusegun Ekanade

**Published by the**  
Department of Geography, Osun State University,  
State of Osun, Nigeria

## EDITORIAL BOARD

**Chairman:**

Dr. R. A. Asiyanbola

**Editor-in-Chief:**

Prof. Olusegun Ekanade

**Members:**

Dr. Samuel Yakubu

Dr. K. J. Samuel

S. A. Adeniyi

A. O. Olusola

O.S. Durowoju

**Business Manager:**

Dr. M.O. Obidiya

**Editorial Advisers:**

Prof. A.S. Gbadegesin -

Prof. C.O. Olatubara -

Prof. F.A. Adesina -

Prof. A.S. Aguda -

Dr. M.O. Olawole -

Prof. T.E. Ologunorisa -

Prof. F.I. Afolabi -

Prof. W.O. Adebayo -

Prof. O.O. Oyesiku -

Prof. B.A. Badejo -

Prof. P.O. Phil-Eze -

Prof. E.F. Ogunbodede -

Prof. L. T.Ajibade -

Prof. A. Olusanya -

Dr. S.R. Akinola -

Dr. O.C. Nwoke -

University of Ibadan

University of Ibadan

Obafemi Awolowo University

Obafemi Awolowo University

Obafemi Awolowo University

Federal University of Technology, Akure

Ekiti State University

Ekiti State University

Olabisi Onabanjo University

Olabisi Onabanjo University

University of Nigeria

Adekunle Ajasin University

University of Ilorin

Osun State University

Osun State University

Osun State University

# CONTENTS

|                                                                                                                                                                                                                |     |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|
| Editorial Board                                                                                                                                                                                                | ii  |
| Contents                                                                                                                                                                                                       | iii |
| Assessment of Changes in Soil Properties Under Different Landuses in Part of Biu Plateau, Borno, Nigeria<br><i>R.D. Abu, E. Yakubu, G.O. Abu, and H.U. Agbeba</i>                                              | 1   |
| Utilization of Primary Health Care Services by Mothers for Maternal and Child Health Services in Rural Niger State, Nigeria<br><i>M. Abubakar and S.A. Abubakar</i>                                            | 9   |
| An Assessment of Road Transport Safety Standardization Scheme Influence on Freight Transport Crashes in Lagos Metropolis, Nigeria<br><i>H.O. Adebayo, C.O. Akanni, M.O. Solanke, B.A. Raji, and K. Olaleye</i> | 25  |
| Street Encroachment by the Informal Metal Container Operators in Osogbo, Osun State, Nigeria<br><i>S.B. Adedotun, D.S. Ogundahunsi and D.O. Adedotun</i>                                                       | 36  |
| An Assessment of the Potential of Gidan Dan Hausa Museum for Sustainable Tourism Development in Kano State, Nigeria<br><i>A.S. Muhammad</i>                                                                    | 43  |
| Quality of Urban Open Spaces in a Southwestern Nigerian City<br><i>A. Ajayi and D. Amole</i>                                                                                                                   | 52  |
| Landowner and Land Fragmentation in Kano Metropolis, Nigeria<br><i>I.M. Dankani</i>                                                                                                                            | 64  |
| Assessment of Climate Variability Perception Among Farmers in Keffi Local Government Area, Nasarawa State, Nigeria<br><i>J.I. Magaji, I.M. Anzaku, A.T. Ogah, A. Mohammed, A.G. Idoko</i>                      | 74  |
| Social and Medical Implications of ICT in Tourism and Hospitality Industry in Nigeria<br><i>M.O. Lawal, Y.K. Irehue and T.Y. Adeyinka</i>                                                                      | 83  |
| Assessment of Urban Landuse Expansion on Agricultural Land in Kwadon District of Yamaltu-Deba Local Government Area, Gombe State<br><i>L.A. Mbaya</i>                                                          | 92  |

|                                                                                                                                                                                            |     |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|
| Spatial Analysis of Adolescents' Accessibility to Contraceptives in Osun State, Nigeria<br><i>M.O. Obidiya</i>                                                                             | 102 |
| Telematics (ICT) Usage in Nigerian Railway Operations: A Case Study of Iddo – Ijoko<br>Route<br><i>F.R. Ohaegbue, I.O. Longe, E.E. Ege and S.I. Oni</i>                                    | 111 |
| Remote Sensing/GIS Software Utilization and Challenges – An Exploratory Study in<br>Nigeria<br><i>R.A. Asiyanbola</i>                                                                      | 118 |
| Anthropogenic Activities, Pressure and Threat on Girei Forest Reserve, Adamawa State,<br>Nigeria<br><i>S. Adamu and A. Musa</i>                                                            | 125 |
| Comparative Analysis of the Characteristics of Rainstorms Over Ibadan, Nigeria<br><i>A. Adediran</i>                                                                                       | 135 |
| The Effects of Heavy Metals Discharged into River Kaduna on the Quality Of Irrigated<br>Farmlands at Kaduna Metropolis, Kaduna State<br><i>Y.M. Abui, B.R. Atiyong, S. Stephen D. Laah</i> | 144 |
| Assessment of Strategic Peri-urban Structure of Ojoo and Sasa Areas of Ibadan, Nigeria<br><i>O. Ekanade, S. Yakubu and D. A. Yakubu</i>                                                    | 153 |
| Socio-economic Impact of Small-scale Industries in Ogbomoso Township, Oyo State,<br>Nigeria<br><i>R.B. Ibrahim, M.A. Adetunji and C.O. Ajala</i>                                           | 163 |
| * Risk Assessment of Toxicity Potential for Criteria Ambient Air Pollutants In Selected<br>Areas of Lagos State, Nigeria<br><i>K.O. Laro and U.A. Raheem</i>                               | 175 |

# RISK ASSESSMENT OF TOXICITY POTENTIAL FOR CRITERIA AMBIENT AIR POLLUTANTS IN SELECTED AREAS OF LAGOS STATE, NIGERIA

K.O. Laro and U.A. Raheem

Department of Geography & Environmental Management, University of Ilorin, Ilorin, Nigeria  
Corresponding author's e-mail: laro.ko@unilorin.edu.ng

## Abstract

The quality of ambient air is determined by the extent of pollution of the environment. The United States Environmental Protection Agency identifies six pollutants namely particulate matter, ground level ozone, carbon monoxide, sulphur oxides, nitrogen oxides and lead exposure as toxic and injurious because of their potentials to cause or trigger various human health problems. This study examines the toxicity potential of criteria ambient air pollutants in selected areas of Lagos State, Nigeria. The data for the study include five-year data of criteria pollutants for selected areas of Lagos and World Health Organization (WHO) guideline values of criteria pollutants. Data analysis adopted the use of Toxicity Potential Model for the calculation of the toxicity potential of the criteria pollutants. GIS procedure was employed to produce a map showing the spatial pattern and hotspot areas of criteria pollutants. Criteria pollutants in the study area were also compared with the WHO standards with a view to determining areas of excessive pollution and toxicity potential in the area. The study reveals that  $PM_{2.5}$ ,  $PM_{10}$ ,  $SO_2$  and  $NO_2$  in Lagos areas clearly exceeded the benchmark set by WHO. Furthermore, CO and  $O_3$  are well below the WHO standards. The toxicity potential values for  $PM_{2.5}$ ,  $PM_{10}$ ,  $SO_2$  and  $NO_2$  in the area of study are greater than unity while the values for CO and  $O_3$  are lesser than unity which indicates that such concentration has a tendency of causing harm to people and hence should be avoided. The study recommends that governments, and other stakeholders in health should harmonize efforts, resources and ideas towards effective planning, monitoring, and provision of facilities that could control and ameliorate the presence of these pollutants to which urban residents are exposed thereby reducing the health effects from such exposures.

**Keywords:** Ambient air, Pollution, Health, Toxicity, Criteria pollutants

## Introduction

The quality of ambient air is determined by the extent of pollution of the environment. Pollution on the whole is caused principally by human activities, though it can also be a natural process (Dara, 2004). The United States Environmental Protection Agency (USEPA, 2012) classified air pollutants into two groups: criteria and hazardous air pollutants. The criteria pollutants are those that can have adverse effect on health and the environment. They include particulate matter, oxides of nitrogen, oxides of sulphur, carbon (II) Oxide and lead. Criteria pollutants are toxic and injurious to health. Apart from their natural presence in air, they can be introduced into the air by various types of

anthropogenic activities such as incomplete combustion from engines, bush-burning, electricity generating plants, decaying of accumulated organic and domestic wastes, and other industrial processes (Raheem and Amole, 2015).

Air quality can be a critical reflection of the ambient atmospheric pollution, relative to the potential to inflict harm on the environments (WHO, 2002). Air is said to be polluted when there is 'the presence in the outdoor atmosphere of one or more contaminants such as dust, fumes, gas, mist, odour, smoke, or vapour in such a characteristics and duration as to make them actually or potentially injurious to human, plant or animal life, or property or which interferes with the comfortable enjoyment of life and

property' (World Bank, 1978). Human health is threatened with diseases and early mortality particularly in emerging economies facing rapid industrialization and urbanization like Lagos. Epidemiological studies of occupational diseases on the working population had shown negative effects of the environment on people working in a contaminated environment over a lifetime of employment (The MACBETH project 1999, and Samoli, E. 2003). According to WHO (2014), by reducing air pollution levels, countries can reduce the burden of disease from stroke, heart disease, lung cancer, and both chronic and acute respiratory diseases, including asthma. Moreover, reduction in outdoor air pollution also reduces emissions of CO<sub>2</sub> and short-lived climate pollutants such as black carbon particles and methane, thus contributing to the near and long-term mitigation of climate change (WHO, 2014).

Air quality has a strong link to human health and wellbeing as shown in many studies (Zhu *et al.* 2002, Parakash, *et al.* 2006). The rapid expansion, population increase and transformation of the Lagos city, one of the most industrialized cities in the world have contributed to its present polluted states. Akinola *et al.* (2014) posited that the increasing human activities, especially industrial and vehicular emissions are posing great environmental challenges that have resulted in loss of life and destruction of properties in the state. The overall effects of air pollution on Lagos residents and environments contribute to drag in the efforts for its most sustainable programs (Akinola *et al.* 2014).

### Statement of the problem

According to Brunekreef (2005), ambient particulate matter levels in cities of developing countries including Nigeria are generally much higher than in developed countries because of dispersed heating with small-scale solid fuel use, uncontrolled industrial emissions, and the large numbers of non-catalyst two-stroke engine vehicles. Ediagbonya *et al.* (2012) posited that atmospheric environmental problems, had previously received scanty attention in Nigeria but have become a subject of increasing importance. Air pollution is a major threat to human life and most people inhale pollutants while at home or commuting to work irrespective of the mode of transportation (Ekpenyong *et al.* 2012). Depending on the dose and the exposure time, these pollutant gases have the

potential to cause far reaching adverse health effects in man, but principally affect the respiratory and cardiovascular systems.

The World Health Organization (2002), argued that about 2.4 million people worldwide (including about 93,700 Nigerians) die each year from causes directly attributable to air pollution. Akinola *et al.* (2014) posited that studies have shown that in Lagos state Vehicular Emissions are the highest point source contributors of carbon dioxide (CO<sub>2</sub>) into the environment, followed by the manufacturing industries. Unending traffic jams in Lagos metropolis results in commuters spending several unproductive hours in traffic and increased avoidable emissions of CO<sub>2</sub> and other pollutants. A study conducted by the Lagos Metropolitan Transport Management Authority (LAMATA) on air quality between 2003 and 2007 indicated that vehicular emission contribute approximately 43% ambient air pollution in Lagos. The continuous expansion, population increase and transformation of the Lagos city have contributed to its present polluted states. However, assessment of the effect of air pollution in developing countries is difficult because of a lack of cohesive air quality policies in combination with poor environmental monitoring and a paucity of disease surveillance data (Briggs, 2003). Therefore, this study becomes imperative in examining the hotspot areas and toxicity potential for criteria ambient air pollutants in selected area of Lagos State.

### Study Area

Lagos State is located between latitudes 6°.35N to 6.58°N and longitudes 3°.45'E to 3.75°E of the Greenwich meridian in the south western part of Nigeria. The state has a tropical wet and dry climate with an all year-round precipitation in many parts of the state. Wet season is characterised by a double maximum of rainfall usually from March to July and the other in late August to early September. A dry spell may occur from late September to early November. The annual mean rainfall is between 1381.7mm and 2733.4mm in recent time from one location to another. The maximum temperature ranges between 29°C - 34°C, the lowest being in the month of July and the highest in February and a minimum temperature varies between 24°C -28°C. The relative humidity varies seasonally with an average of 70% throughout the year.

The vegetation of the study area is made up of two types namely; swamp forest of the coastal belt and

dry lowland rain forest. The swamp forest is a combination of mangrove forest and coastal vegetation developed under the brackish conditions of the coastal areas, swamp fresh water lagoons and estuaries. Lying to the north of the swamp forest is the lowland (tropical) rain forest zone and stretches from Ikeja through Ikorodu. The economic valuable trees such as teak, tripochiton, seletrocylon (arere), bancia diderrichil (opepe) and terminahia (idigbo) can be found in some parts of the study area. Lagos State occupies an area of 3,577 square kilometres, which represents 0.4% of Nigeria landmass with a marine shoreline of about 180km extending inland to a maximum distance of about 32km. Lagos is the most populous city in Nigeria, the second fastest-growing city in Africa and the seventh in the world ([www.lagosstate.gov.ng](http://www.lagosstate.gov.ng)). The NPC (2016) estimates of the population of Lagos state was 21 million, making Lagos the largest city in Africa.

The study area accounts for over 60% of the federation's total industrial investment and the largest concentrations of industries can be found in Ikeja, Alimosho and Kosofe local government areas ([www.lagosstate.gov.ng](http://www.lagosstate.gov.ng)). Other specific locations of numerous industries include Apapa, Surulere, Shomolu, Mushin, Oshodi-Isolo, Agege, Amuwo Odofin, and Ikorodu among others. See fig. 1 for the map of the study area.

## Aim and objectives

The aim of the study is to identify the hotspot areas and toxicity potential for criteria ambient air pollutants in selected area of Lagos State. To achieve this aim, the specific objectives are to: compare criteria pollutants in Lagos area with the WHO standard for the pollutants and identify hotspot areas and toxicity potential for criteria pollutants based on deviations from the WHO benchmark.

## Materials and Methods

### Data required and sources

The US Environmental Protection Agency (USEPA, 2012) identifies six common air pollutants also known as criteria pollutants. These are particulate matter ( $PM_{2.5}$ ,  $PM_{10}$ ), ground level ozone, carbon monoxide, sulphur oxides, nitrogen oxides and lead. These pollutants are critical because of their implications for human health and the environment. Out of the six, particulate matter and ozone are the most widespread health threats because of their association with Asthma, Chronic Obstructive Pulmonary Disease (COPD) and respiratory diseases as prominent air pollution related diseases (WHO, 2014).




Figure 1. Lagos State showing sample areas with Nigeria as inset

Based on the foregoing the data required and sources of data for this study are both primary and secondary. The primary sources includes the coordinates of the sample sites. The secondary data includes five (5) years data of criteria pollutants, the pollutants are Particulate matter ( $PM_{2.5}$ ,  $PM_{10}$ ), Ground level ( $O_3$ )

Nitrogen dioxide ( $NO_2$ ), sulphur dioxide ( $SO_2$ ), and carbon dioxide ( $CO$ ), WHO Guidelines for  $PM_{2.5}$ ,  $PM_{10}$ ,  $SO_2$ ,  $NO_2$ ,  $CO$  and Ground level  $O_3$ , maps, and statistical data obtained from WHO published works.

**Table 1:** Showing Study Sites Distribution within the Study Area.

| LANDUSE TYPE | LOCATIONS/SITES                               |
|--------------|-----------------------------------------------|
| RESIDENTIAL  | Agege Housing Estate; Oko-oba                 |
|              | Ikeja G.R.A: SoboArobiudu street.             |
|              | Magodo Estate: CMD Road, Shangisha            |
|              | Ogba Staff Quarters; Ogba                     |
|              | Secretariat; Alausa: front of Skye bank       |
| HIGH TRAFFIC | Maryland Junction, Ikeja                      |
|              | General Hospital: Frontage of LASUTH          |
|              | Ojodu-Berger: Berger Roundabout               |
|              | Iyana-Ipaja: Under Bridge                     |
|              | Oshodi B/Stop: Oshodi                         |
| INDUSTRIAL   | Specomil: Inside Ikeja Industrial Estate      |
|              | Wemabod: Adjacent Guinness Nigeria PLC office |
|              | Wempco: Industrial Estate Ogba                |
|              | Ize-Iyamu                                     |
|              | Matori                                        |

Source: Author's Fieldwork, 2016




**Figure 2:** Sampling Sites in the Study Area

Source: Author's Fieldwork, 2016

### Methods of Data Collection

Study sites were selected with a view to giving a holistic representation of air quality status of the study area. The study area was categorised into three (3) base on the predominant landuse types. Thus, Lagos was divided into high traffic, industrial and residential areas. Residential areas were defined as landuse types in which housing predominates. In the light of the above, State Government staff quarters and Estates owned by private agencies were considered. The high traffic areas are the major road intersections where high volumes of traffic are experienced during the working hours of the day while areas with high concentration of industries were considered as industrial areas. It must be noted that this categorisation is not water tight as these characteristics may occur in all areas with varying intensities (Laro, 2016)

Geographic coordinates: Direct measurement of coordinates of fifteen (15) sample sites from the field with the aid of the global positioning system (GPS).

Data on Criteria pollutants: The data on criteria pollutants was obtained from Ministry of Environment, Lagos State. The data are particulate matter ( $PM_{2.5}$  and  $PM_{10}$ ), Ground level Ozone ( $O_3$ ), Nitrogen dioxide ( $NO_2$ ), Sulphur dioxide ( $SO_2$ ) and Carbon monoxide (CO).

Guideline value for criteria pollutants: The data was sourced from the WHO air quality guideline book. The data are annual, daily and 8 hours mean of  $PM_{2.5}$ ,  $PM_{10}$ ,  $SO_2$ ,  $NO_2$ , CO and ground level ( $O_3$ ).

### Methods of Data Analysis

Descriptive statistics was employed to compare Criteria Pollutants in Lagos area with the WHO standard. For the purpose of this study, WHO Values which are presented in  $\mu g/m^3$  are converted to  $mg/m^3$  using gas detection calculator ([www.gasdetectioncalculator.com](http://www.gasdetectioncalculator.com)). This was because the data on criteria pollutants collected from Lagos state Ministry of Environment were presented in  $mg/m^3$ . Hotspot areas and toxicity potential for Criteria Pollutants were obtained as deviations from the WHO Guidelines and Standard defining the permissible level of each criteria pollutants

Toxicity Potential is expressed as the ratio of measured ambient pollutants concentration to the statutory limit of ambient concentration (Sonibare *et al.* 2005). It is useful in assessing the deleterious effects of air pollutants emissions on human health.

It was computed using Eqn (1) taking into consideration the ambient air quality standards of World Health Organisation (WHO) as reported in WHO Air quality guidelines (2005).

The toxicity potential of the pollutants measured was computed using the following equation:

$$TP_i = \frac{Ci(t)}{Sli(t)} \dots \dots \dots \text{eq. (1)}$$

Where TP represents the toxicity potential of pollutant 'i'

This equation takes into consideration the ambient air quality standard of WHO as reported in WHO air quality guidelines (2005)

In the same vein, the use of tables was employed to identify the hotspot areas in ArcGIS environment. The discussion and interpretation of the results are presented in the section below.

## Results and Discussion

### Benchmarking criteria pollutants in Lagos

The WHO (2005) air quality guidelines book presents revised guideline values for the four most common air pollutants - particulate matter ( $PM_{2.5}$  and  $PM_{10}$ ), nitrogen dioxide, sulphur dioxide and ozone based on a recent review of the accumulated scientific evidence. The rationale for selection of each guideline value is supported by a synthesis of information emerging from research on the health effects of each pollutant. Based on the foregoing, these guidelines are apply globally (WHO, 2005). The details of the WHO Air Quality guidelines is as shown in table 2

As shown in Table 3, Ambient  $PM_{2.5}$  and  $PM_{10}$  level in the study area were observed to be higher than the internationally permitted standard of  $0.01 \text{ mg}/\text{m}^3$  and  $0.02 \text{ mg}/\text{m}^3$  respectively over an annual average period (WHO, 2005). The ambient level of carbon monoxide (CO) in the study area were observed to be lower than the statutory standard of 10 ppm. Also,  $PM_{2.5}$  and  $PM_{10}$  level in the study area are higher than the WHO recommended value of 0.02 ppm and 0.01 ppm respectively except for Agege housing estates, which is a residential area where 0.01 ppm was recorded for both pollutants. Ambient ground level ozone ( $O_3$ ) concentration in the study area were

observed to be lower than the WHO statutory standard of 0.05 ppm over an eight (8) hours averaging period.

### Hotspot areas for criteria pollutants and toxicity potentials in Lagos Metropolis

As indicated in Table 4, the hotspot areas are where there was high concentration of criteria pollutants in the area of study. These areas deviate above the recommended WHO standard, the areas were highlighted in 'reds' signifying hotspot areas. The table also reveals that a minimum of two (2) and maximum of four (4) pollutants across different landuse in the study area exceeds the statutory recommended value.

The minimum toxicity potential with WHO standard for PM<sub>2.5</sub> was 8 computed for Agege housing estate, a residential area, while the maximum was 33 in Ojodu Berger, a high traffic area (see Table 5). The minimum toxicity potential for PM<sub>10</sub> was 7 in Ogba Staff quarters which is a residential area, while the

maximum was 23 in Iyana-Ipaja, a high traffic area. Also, the minimum toxicity potential for CO was 0.01 in Ogba Staff quarters and Secretariat, a residential area, while the maximum was 0.18 in Ojodu Berger which was a high traffic area. The minimum toxicity potential for NO<sub>2</sub> was 0.5 ppm in Agege housing estate, a residential area, while the maximum was 17.5 ppm in Ojodu Berger a high traffic area. The minimum toxicity potential for SO<sub>2</sub> was 1 in Agege, a residential area, while the maximum is 43 in Ojodu Berger a high traffic area. Minimum toxicity potential for O<sub>3</sub> was 0.08 in Ikeja GRA, a residential area, while the maximum is 0.6 in General Hospital and Iyana-Ipaja a high traffic areas, and Ize-Iyamu and Matori, an industrial areas. It follows therefore that high traffic areas recorded the highest values of criteria pollutants, followed by industrial and residential areas for instance 33, 19 mg/m<sup>3</sup> and 17.5, 43ppm was recorded for PM<sub>2.5</sub>, PM<sub>10</sub> and NO<sub>2</sub>, SO<sub>2</sub> respectively in Ojodu Berger. By implications, High traffic and industrial areas as indicated in Table 5 has the highest toxicity level of criteria pollutants.

**Table 2:** Conversion of WHO Guideline values for pollutants from microgram ( $\mu\text{g}/\text{m}^3$ ) to milligram ( $\text{mg}/\text{m}^3$ )

| Pollutants                              | Guideline value in ( $\mu\text{g}/\text{m}^3$ / ppm) | Value in ( $\text{mg}/\text{m}^3$ / ppm) |
|-----------------------------------------|------------------------------------------------------|------------------------------------------|
| Particulate matter (PM <sub>2.5</sub> ) | 10 $\mu\text{g}/\text{m}^3$ (annual mean)            | 0.01 $\text{mg}/\text{m}^3$              |
| Particulate matter (PM <sub>10</sub> )  | 20 $\mu\text{g}/\text{m}^3$ (annual mean)            | 0.02 $\text{mg}/\text{m}^3$              |
| Nitrogen dioxide (NO <sub>2</sub> )     | 40 $\mu\text{g}/\text{m}^3$ (annual mean)            | 0.02 ppm                                 |
| Sulphur dioxide (SO <sub>2</sub> )      | 20 $\mu\text{g}/\text{m}^3$ (daily mean)             | 0.02 ppm                                 |
| Ozone (O <sub>3</sub> )                 | 100 $\mu\text{g}/\text{m}^3$ (8hr mean)              | 0.05 ppm                                 |
| Carbon monoxide (CO)                    | 10 ppm (8hr mean)                                    | 10 ppm                                   |

\* Conversion factor: [gasdetectioncalculator.com](http://gasdetectioncalculator.com)

Source: Authors computation, 2016.

**Table 3:** Criteria Pollutants and International Standards

| Landuse type             |                | PM <sub>2.5</sub><br>( $\text{mg}/\text{m}^3$ ) | PM <sub>10</sub><br>( $\text{mg}/\text{m}^3$ ) | CO<br>(ppm) | NO <sub>2</sub><br>(ppm) | SO <sub>2</sub><br>(ppm) | O <sub>3</sub><br>(ppm) |
|--------------------------|----------------|-------------------------------------------------|------------------------------------------------|-------------|--------------------------|--------------------------|-------------------------|
| Residential<br>Areas     | Ogba S/Qtrs    | 0.12                                            | 0.14                                           | 0.12        | 0.03                     | 0.03                     | 0.01                    |
|                          | Secretariat    | 0.14                                            | 0.10                                           | 0.14        | 0.03                     | 0.02                     | 0.01                    |
|                          | Magodo         | 0.14                                            | 0.13                                           | 0.24        | 0.06                     | 0.05                     | 0.01                    |
|                          | Ikeja GRA      | 0.16                                            | 0.14                                           | 0.20        | 0.04                     | 0.04                     | 0.004                   |
|                          | Agege H/Estate | 0.08                                            | 0.09                                           | 0.09        | 0.01                     | 0.01                     | 0.01                    |
|                          | Maryland       | 0.22                                            | 0.11                                           | 0.85        | 0.22                     | 0.30                     | 0.02                    |
| High<br>Traffic<br>Areas | Oshodi         | 0.26                                            | 0.27                                           | 1.10        | 0.26                     | 0.36                     | 0.02                    |
|                          | Ojodu Berger   | 0.33                                            | 0.38                                           | 1.76        | 0.35                     | 0.43                     | 0.02                    |
|                          | Gen.Hospital   | 0.25                                            | 0.19                                           | 0.87        | 0.23                     | 0.34                     | 0.03                    |
|                          | Iyana-Ipaja    | 0.30                                            | 0.46                                           | 1.16        | 0.40                     | 0.38                     | 0.03                    |
| Industrial<br>Areas      | Specomil       | 0.26                                            | 0.28                                           | 0.49        | 0.06                     | 0.09                     | 0.02                    |
|                          | Wernabod       | 0.17                                            | 0.15                                           | 0.37        | 0.04                     | 0.07                     | 0.02                    |
|                          | Wempco         | 0.17                                            | 0.19                                           | 0.35        | 0.09                     | 0.11                     | 0.02                    |
|                          | Ize-Iyamu      | 0.23                                            | 0.26                                           | 0.73        | 0.13                     | 0.12                     | 0.03                    |
| Int.<br>Standard         | Matori         | 0.23                                            | 0.19                                           | 0.70        | 0.07                     | 0.09                     | 0.03                    |
|                          | WHO            | 0.01                                            | 0.02                                           | 10          | 0.02                     | 0.01                     | 0.05                    |

Source: Author's Computation, 2016

**Table 4:** Hotspots in the Three Land use Areas

| Locations          |                | PM <sub>2.5</sub><br>(mg/m <sup>3</sup> ) | PM <sub>10</sub><br>(mg/m <sup>3</sup> ) | CO<br>(ppm) | NO <sub>2</sub><br>(ppm) | SO <sub>2</sub><br>(ppm) | O <sub>3</sub><br>(ppm) |
|--------------------|----------------|-------------------------------------------|------------------------------------------|-------------|--------------------------|--------------------------|-------------------------|
| Residential Areas  | Ogba S/Qrtrs   | 0.12                                      | 0.14                                     | 0.12        | 0.03                     | 0.03                     | 0.01                    |
|                    | Secretariat    | 0.14                                      | 0.10                                     | 0.14        | 0.03                     | 0.02                     | 0.01                    |
|                    | Magodo         | 0.14                                      | 0.13                                     | 0.24        | 0.06                     | 0.05                     | 0.01                    |
|                    | Ikeja GRA      | 0.16                                      | 0.14                                     | 0.20        | 0.04                     | 0.04                     | 0.004                   |
|                    | Agege H/Estate | 0.08                                      | 0.09                                     | 0.09        | 0.01                     | 0.01                     | 0.01                    |
|                    | Maryland       | 0.22                                      | 0.11                                     | 0.85        | 0.22                     | 0.30                     | 0.02                    |
| High Traffic Areas | Oshodi         | 0.26                                      | 0.27                                     | 1.10        | 0.26                     | 0.36                     | 0.02                    |
|                    | Ojodu Berger   | 0.33                                      | 0.38                                     | 1.76        | 0.35                     | 0.43                     | 0.02                    |
|                    | Gen.Hospital   | 0.25                                      | 0.19                                     | 0.87        | 0.23                     | 0.34                     | 0.03                    |
|                    | Iyana-Ipaja    | 0.30                                      | 0.46                                     | 1.16        | 0.4                      | 0.38                     | 0.03                    |
| Industrial Areas   | Specomil       | 0.26                                      | 0.28                                     | 0.49        | 0.06                     | 0.09                     | 0.02                    |
|                    | Wemabod        | 0.17                                      | 0.15                                     | 0.37        | 0.04                     | 0.07                     | 0.02                    |
|                    | Wempco         | 0.17                                      | 0.19                                     | 0.35        | 0.09                     | 0.11                     | 0.02                    |
|                    | Ize-Iyamu      | 0.23                                      | 0.26                                     | 0.73        | 0.13                     | 0.12                     | 0.03                    |
| Int. Standards     | Matori         | 0.23                                      | 0.19                                     | 0.70        | 0.07                     | 0.09                     | 0.03                    |
|                    | WHO            | 0.01                                      | 0.02                                     | 10          | 0.02                     | 0.01                     | 0.05                    |

\*Reds signifies hotspot areas

Source: Authors Computation, 2016.

**Table 5:** Computed Toxicity Potential of the Criteria Pollutants

| Locations          |                | PM <sub>2.5</sub><br>(mg/m <sup>3</sup> ) | PM <sub>10</sub><br>(mg/m <sup>3</sup> ) | CO<br>(ppm)       | NO <sub>2</sub><br>(ppm) | SO <sub>2</sub><br>(ppm) | O <sub>3</sub><br>(ppm) |
|--------------------|----------------|-------------------------------------------|------------------------------------------|-------------------|--------------------------|--------------------------|-------------------------|
|                    |                | TP <sub>WHO</sub>                         | TP <sub>WHO</sub>                        | TP <sub>WHO</sub> | TP <sub>WHO</sub>        | TP <sub>WHO</sub>        | TP <sub>WHO</sub>       |
| Residential Areas  | Ogba S/Qrtrs   | 12                                        | 7                                        | 0.01              | 1.5                      | 3                        | 0.2                     |
|                    | Secretariat    | 14                                        | 5                                        | 0.01              | 1.5                      | 2                        | 0.2                     |
|                    | Magodo         | 14                                        | 6.5                                      | 0.02              | 3                        | 5                        | 0.2                     |
|                    | Ikeja GRA      | 16                                        | 7                                        | 0.02              | 2                        | 4                        | 0.08                    |
|                    | Agege H/Estate | 8                                         | 4.5                                      | 0.09              | 0.5                      | 1                        | 0.2                     |
|                    | Maryland       | 22                                        | 5.5                                      | 0.09              | 11                       | 30                       | 0.4                     |
| High Traffic Areas | Oshodi         | 26                                        | 13.5                                     | 0.11              | 13                       | 36                       | 0.4                     |
|                    | Ojodu Berger   | 33                                        | 19                                       | 0.18              | 17.5                     | 43                       | 0.4                     |
|                    | Gen. Hosp.     | 25                                        | 9.5                                      | 0.09              | 11.5                     | 34                       | 0.6                     |
|                    | Iyana-Ipaja    | 30                                        | 23                                       | 0.12              | 20                       | 38                       | 0.6                     |
| Industrial Areas   | Specomil       | 26                                        | 14                                       | 0.05              | 3                        | 9                        | 0.4                     |
|                    | Wemabod        | 17                                        | 7.5                                      | 0.04              | 2                        | 7                        | 0.4                     |
|                    | Wempco         | 17                                        | 9.5                                      | 0.04              | 4.5                      | 11                       | 0.4                     |
|                    | Ize-Iyamu      | 23                                        | 13                                       | 0.07              | 6.5                      | 12                       | 0.6                     |
|                    | Matori         | 23                                        | 9.5                                      | 0.07              | 3.5                      | 9                        | 0.6                     |

\*Reds signifies toxicity potential values above unity

Source: Author's Computation, 2016

It must be noted that toxicity potential values greater than unity indicates that such concentration has a tendency of causing harm to people that are exposed to it in such an environment and hence should be avoided. The toxicity potentials values highlighted in Table 5 are areas with concentration of criteria pollutants that has a potential for harmful effects to people. Fig. 3 shows the map of the hotspot areas.

Subsequently, Figure 3 shows the hotspot areas with high toxicity potential, these areas pose a serious health threat to the inhabitants around the areas and

major road user's. For instance, Burnnet *et al* (1998) found a 4.3% increase in all-cause mortality for an increase in 24-hour average NO<sub>2</sub> levels 47µg/m<sup>3</sup>. Dennison *et al* (2002) summarises the health effects of associated with exposure to Ozone as increase in daily Mortality, Respiratory and Cardiovascular diseases such as High blood Pressure, Cough, Phlegm and wheeze.

Long-term exposure to PM<sub>2.5</sub> is associated with an increase in the long-term risk of cardiopulmonary mortality by 6–13% per 10 µg/m<sup>3</sup>

of  $PM_{2.5}$  (Pope, C.A. III, 2002). Susceptible groups with pre-existing lung or heart disease, as well as elderly people and children, are particularly vulnerable. For example, exposure to PM affects lung development in children, including reversible deficits in lung function as well as chronically reduced lung growth rate and a deficit in long-term lung function

(WHO, 2011). Dennison *et al* (2002) concludes that because of the correlations between  $SO_2$  and other contaminants in the air it is difficult to confidently attribute the observed effects in the epidemiological studies to  $SO_2$  alone.




**Figure 3:** Spatial pattern of Hotspot areas of pollutants

Source: Authors Fieldwork, 2016

## Conclusion

Spatial variation in the distribution of Criteria pollutants in Lagos was established. From the findings, it was evident that  $PM_{2.5}$ ,  $PM_{10}$ ,  $SO_2$  and  $NO_2$  in sample areas clearly exceeded the regulatory limit set by WHO especially in the high traffic zones. Furthermore, CO and  $O_3$  are well below the statutory standards. Areas such as Ojodu-berger, Oshodi, Ojota, Ikeja and Iyana-ipaja were classified as high risk areas. Also, the toxicity potential values for  $PM_{2.5}$ ,  $PM_{10}$ ,  $SO_2$  and  $NO_2$  in the area of study are greater than unity while the values for CO and  $O_3$  are lesser than unity. However, toxicity potential values greater than unity indicates that such concentration has a tendency of causing harm to people and hence should be avoided. This was because most of the criteria pollutants exceeded the recommended WHO value.

## Recommendations and Policy Measures

There is need for better air quality in the state in order

to reduce the health effect of air pollutants. These could be achieved, if only the Government could implement the following recommendations that was made based on the findings of this study. Government should create Lagos State air quality health index (AQHI) and report on cumulative health impacts associated with criteria pollutants to be monitored at suitable locations within the state. Enforcement agencies such as Lagos State Traffic Management Agency (LASTMA) and LAMATA should ensure the compliance of Industrial and vehicular emission standard and provision of database for determining air quality in major metropolitan area of the state. There are needs to encourage mass transportations throughout the state, cleaner modes of transportation and more sustainable energy generation in the state. Also, tree planting and greening of the environment should be sustained and improved upon throughout the state for aesthetics and carbon sequestration purposes.

## References

- Akinola, A. K., Suleiman, A. A., Adeleye, Y. B., Anifowose, F.O. & Dauda, R. A. (2014). Air pollution and climate change in Lagos, Nigeria: Needs for proactive approaches to risk management and adaptation. *American Journal of Environmental Sciences*, 10(4), 412-423.
- Briggs, D. (2003). Environmental pollution and the global burden of diseases. *Br Med Bulletin*, 68, 1-24.
- Brunekreef, B. (2005). Out of Africa: *Occupational and Environmental Medicine Journal*, 62(6), 351-2.
- Burnett, R. T., Cakmak, S. & Brook, J., R. (1998). The effect of urban ambient air pollution mix on daily mortality rates in eleven Canadian cities. *Canadian Journal of Public Health*, 89(3), 152-6.
- Dara, S. S. (2004). *A textbook of Environmental Chemistry and Pollution Control*. New Delhi, S, Chand and Company Limited
- Dennison, L., Rolfe, K. & Graham, B. (2002). Review of the ambient air quality guidelines – health effect of five common air contaminants and recommended protective ranges. Ministry for the Environment Technical Report No. 12
- Ediagbonya, T. F., Ukpebor, E. E., Okieimen, F. E. & Okungbowa, G. E. (2012). Comparative study of TSP, inhalable particles and respirable particles in urban and rural areas in Niger Delta region of Nigeria. *Greener Journal of Physical Sciences*, 2(3), 89-96
- Ekpenyong, C. E., Ettebong, E. O., Akpan, E. E., Samson, T. K. & Daniel, N. E. (2012). Urban city transportation mode and respiratory health effect of air pollution: A cross-sectional study among transit and non-transit workers in Nigeria. *BMJ OPEN*, 2, 5-20
- Laro, K. O. (2016). Ambient air quality and incidence of selected diseases in Lagos, Nigeria. An unpublished MSc thesis, submitted to the department of geography and environmental management, University of Ilorin, Ilorin, Nigeria
- Parakash, E., Gauderman, W. J. & Peters, J. (2006). Childhood asthma and exposure to traffic and nitrogen dioxide, *Epidemiology Journal*, 16(2), 737 – 743.
- Pope, C. A. III (2002). Lung cancer, cardiopulmonary mortality, and long-term exposure to fine particulate air pollution. *Journal of the American Medical Association*, 287(9), 1132-1141.
- Raheem, U. A., & Amole, S. O. (2015). Methane emission from municipal solid waste in Ilorin, Nigeria: A risk analysis for household health.

- Journal of Environmental Science Technology*, Centre for human and urban settlement development. Federal University of Technology, Minna
- Samoli, E. (2003). Acute effects of ambient particulate matter on mortality in Europe and North America: Results from the APHENA Study. *Environmental Health Perspectives*, 116 (11), 1480–1486.
- Sonibare, J. A., Akeredolu, F. A., Osibanjo, O. & Latinwo, I. (2005). Analysis of total suspended particulates from Enamelware Manufacturing Industry, *American Journal of applied science*, 2 (2), 573-578
- The MACBETH Project (1999). Monitoring of Atmospheric Concentrations of Benzene in European Towns and Homes, *European Union LIFE Project 96 ENV / IT070*.
- USEPA (2012). Criteria Air Pollutants. Retrieved from <http://www.epa.gov/air/urbanair> Accessed September 15, 2015.
- World Bank (1978). Air Pollution, Atmospheric air pollution in Industrialised nations, *World Bank Journal*, 2 (4), 12-14
- WHO (2002). Systemic review of health aspects of AQ Europe. An overview of the St. George's project. A systemic review of the epidemiological literature on the short-term health effects of outdoor air pollution, St. George's Hospital, London, United Kingdom
- WHO (2005). Air Quality Guidelines: Global Update. Particulate Matter, Ozone, Nitrogen Dioxide and Sulphur Dioxide. Retrieved from <http://www.euro.who.int/en/health-topics/environment-and-health/Housing-and-health/publications/pre-2009/air-quality-guidelines.-global-update-2005.-particulate-matter,-ozone,-nitrogen-dioxide-and-sulfur-dioxide>. Accessed March 16, 2016.
- WHO (2011). Exposure to Air Pollution (Particulate Matter) In Outdoor Air. Copenhagen, WHO Regional Office for Europe, 2011 (ENHIS Factsheet 3.3) ( \_3.3\_July\_2011.pdf, accessed September 12, 2015.
- WHO (2014). Air quality and health, fact sheet No. 313, WHO Media Centre. Retrieved from <http://www.who.int/mediacentre/factsheets/fs313/en/>. Accessed September 10, 2015
- <http://www.gasdetection.com/calculator/mic2ppm.xls>. Retrieved 25<sup>th</sup> March, 2016
- <http://www.lagosstategov.ng/> retrieved 25<sup>th</sup> March, 2016