

A review of bovine fasciolosis and other trematode infections in Nigeria

N. Elelu^{1,2*} and M.C. Eisler²

¹Faculty of Veterinary Medicine, University of Ilorin, Kwara State, Nigeria:

²University of Bristol, School of Veterinary Science, Langford, Bristol, BS40 5DU, UK

(Received 23 January 2017; Accepted 24 April 2017; First published online 22 May 2017)

Abstract

Trematode infections cause serious economic losses to livestock worldwide. Global production losses due to fasciolosis alone exceed US\$3 billion annually. Many trematode infections are also zoonotic and thus a public health concern. The World Health Organization has estimated that about 56 million people worldwide are infected by at least one zoonotic trematode species, and up to 750 million people are at risk of infection. Fasciolosis caused by the fluke *Fasciola gigantica* is endemic in Nigeria and is one of the most common causes of liver condemnation in abattoirs. Total cattle losses from *Fasciola* infection in Nigeria have been estimated to cost £32.5 million. Other trematode infections of cattle, including paramphistomosis, dicrocoeliasis and schistosomiasis, have all been reported in various parts of Nigeria, with varying prevalence. Most publications on trematode infections are limited to Nigerian local and national journals, with very few international reports. This paper therefore summarized the current data on distribution, control and zoonotic trematode infections in Nigeria and other African countries. We also identified research gaps and made recommendations for future research and areas for funding for policy/planning.

Introduction

Nigeria is one of the four leading livestock producers in Sub-Saharan Africa (Aregheore, 2009), contributing up to 12.7% of the total Nigerian agricultural gross domestic products (CBN, 1999). In the tropics, cattle are generally reared under transhumance husbandry systems with little supplementary feeding, resulting in low productivity and high pre-weaning mortality (Ulayi *et al.*, 2007). Also, acute shortage of feeds during the dry season remains a common occurrence, compelling these animals to graze around water bodies that are often heavily infested with potential intermediate hosts of trematode infections (Ulayi *et al.*, 2007).

Trematode infections of cattle, including fasciolosis (Babalola & Schillhorn van Veen, 1976; Dipeolu *et al.*, 2000), paramphistomosis (Bogatko, 1975), dicrocoeliasis (Schillhorn van Veen *et al.*, 1980; Ulayi *et al.*, 2007) and

schistosomiasis (Pugh *et al.*, 1980; Ndifon *et al.*, 1988) have all been reported in various parts of Nigeria. They are referred to as 'digenetic' because they require at least two different kinds of hosts for their full development. The first are the intermediate hosts of mollusc species (slugs, snails or shellfish) while the final hosts are vertebrates (Taylor, 1964). Mixed infection with more than one trematode species is a common occurrence in animals, due to similarities in life cycles. *Fasciola* infection has been reported to occur concurrently with *Schistosoma* and paramphistome species (Schillhorn van Veen *et al.*, 1980; Yabe *et al.*, 2008). It is also reported to occur concurrently with *Dicrocoelium* species (Schillhorn van Veen *et al.*, 1980; Ulayi *et al.*, 2007). Infection with trematode parasites can lead to severe losses to farmers, hence affecting sustainability of food production (Alvarez Rojas *et al.*, 2014).

Fasciolosis, which is also referred to as distomatosis or liver-fluke disease, is a parasitic disease caused by the trematode of the genus *Fasciola*. The most important species are *F. hepatica* and *F. gigantica*, found in the temperate and tropical regions of the world, respectively. The

*Email: nusyelelu@yahoo.com

geographical distribution of *Fasciola* species is determined by the distribution of the snail intermediate hosts (Boray, 1985). The temperate disease occurs in Europe, America and Oceania, while tropical fasciolosis is common in Africa, the Indian sub-continent, Central and South-East Asia, and other subtropical and tropical areas of the world (FAO, 1993). Both species overlap in many areas of Africa and Asia (Walker *et al.*, 2008). Adults of each species can occur concurrently in the same animal host, either because of local overlap or because of livestock movement (Mas-Coma *et al.*, 2009). The two *Fasciola* species may thus interbreed, resulting in hybrid species (Agatsuma *et al.*, 2000; Ashrafi *et al.*, 2006). They are hermaphrodites and are found in the bile ducts of a large number of herbivorous ruminants, equines, pigs and rabbits (FAO, 1993), and wild animals (Hammond, 1972). Humans are also suitable hosts (Boray, 1969). The intermediate hosts of *F. gigantica* are true water snails belonging to the phylum Mollusca, class Gastropoda and subclass Pulmonata (Wright, 1971). They belong to the family Lymnaeidae and super species *Lymnaea* (*Radix*) *auricularia sensu lato*.

Paramphistomes Fischöder 1901, also known as rumen flukes, are gastrointestinal trematodes belonging to the family of Paramphistomatidae (Soulsby, 1982). These are conically shaped flukes measuring 5–12 mm × 2–4 mm. The adults' predilection sites are the rumen and reticulum of ruminants while the immature parasites are found in the small intestines and abomasum (Rojo-Vázquez *et al.*, 2012). Snails of the families Planorbidae, Bulinidae and Lymnaeidae act as intermediate hosts (Soulsby, 1982; Castro-Trejo *et al.*, 1990). They are largely non-pathogenic but clinical outbreaks have been reported to occur. The most important species in Africa is *Paramphistomum microbothrium* (Dinnik, 1964). Others are *Paramphistomum cervi* (European Environment Agency, 2004), *Paramphistomum ichikawai* (Australasia) (FAO, 1993) and *Paramphistomum daubneyi* (first described in Kenya and common in Europe) (Abrous *et al.*, 1996). Recently, there have been reports of an increasing number of cases of rumen flukes, identified mainly as *Calicophoron daubneyi*, in cattle and sheep in the Republic of Ireland (Zintl *et al.*, 2014; Toolan *et al.*, 2015). The adult paramphistomes are regarded as commensals in the rumen, as heavy infections are tolerated without causing any damage to the rumen (Dinnik, 1964), although immature parasites in the small intestine cause clinical disease (Aiello, 1998).

Dicrocoelium species, commonly referred to as lancet flukes, are found in the bile duct of domestic and wild ruminants (Otranto & Traversa, 2002; Taylor *et al.*, 2007). Other animals, such as rabbits, pigs, dogs, horses and humans, can also be infected (Rojo-Vázquez *et al.*, 2012). There are several species, with *D. dendriticum* (Otranto *et al.*, 2007) having the widest distribution worldwide while *D. hospes* is common in Africa (Aiello, 1998). Other species are *D. chinensis* and *D. suppereri* (Rojo-Vázquez *et al.*, 2012). Two intermediate hosts are required. The first intermediate host is a terrestrial snail of the genus *Limicolaria*, while the second intermediate hosts are brown ants of the genus *Formica* (Soulsby, 1982).

Schistosoma species are elongate, unisexual trematodes, commonly referred to as blood flukes and widely distributed throughout Africa, the Middle East, Asia and some Mediterranean countries (Soulsby, 1982). They are found

in the blood vessels, such as the portal, mesenteric and intestinal veins, of domestic animals (Aiello, 1998). The species, *Schistosoma matteei* and *S. bovis* are common in Africa, the Middle East and southern Europe (Vercruysse & Gabriel, 2005; Taylor *et al.*, 2007). *Schistosoma bovis* is the most pathogenic in animals in Africa (Vercruysse & Gabriel, 2005). It occurs in the portal and mesenteric vessels of cattle, sheep and goats. It is similar to the human parasite *S. haematobium* (Soulsby, 1982). *Schistosoma matteei* has also been reported in sporadic human infections. Other African species in ruminants include *S. curassoni* (common in western Africa), *S. margrebowiei* and *S. leiperi*. In Asia, *S. spindale*, *S. nasale* (nasal worm), *S. indicum*, *S. incognitum* and *S. japonicum* are zoonotic in the Far East (De Bont & Vercruysse, 1997). Aquatic snails such as *Bulinus*, *Physopsis*, *Oncomelania*, *Lymnaea* and *Indoplanorbis* act as the intermediate hosts (Taylor *et al.*, 2007).

This review aims at describing the current status of bovine fasciolosis and other trematode infections in Nigeria and surrounding African countries, in relation to economic losses, distribution, molecular studies, control strategies and human infections. Data were collected from publications obtained from an online search as well as local national journals. Bovine fasciolosis reported in the literature across different states of Nigeria were mapped by using QGIS® software v2.12 (Lyon, France).

Economic losses from trematode infections

Tropical fasciolosis caused by *Fasciola gigantica* is regarded as one of the most important single helminth infections of ruminants in Asia and Africa (Boray, 1985; Fabiyi, 1987). Economic losses from fasciolosis are often difficult to estimate and may result directly from increased liver condemnation or indirectly from decreased livestock productivity (Taylor, 1964). Although direct losses are easier to measure, indirect losses are considered to be far more economically important (Kaplan, 2001). With the total cattle population in Africa estimated at 201 million animals, an annual loss of about US\$840 million from fasciolosis infection is predicted (Spithill *et al.*, 1999), but this cost is likely to have increased significantly in the past 18 years. Production losses in cattle infected with liver flukes often take the form of reduced milk production in dairy cattle and poor feed conversion in beef cattle (Armstrong, 1982). A linear relationship between the burden of adult *F. gigantica* and weight gain of cattle has been described, with infected animals gaining only about half the annual weight compared to controls (Sewell, 1966). The amount of weight loss may also, however, be dependent on age, level of nutrition and intensity of infection (Spithill *et al.*, 1999). Other clinical manifestations are anaemia, reduced fertility and reduced work capacity (Hillyer, 2005), and where there is massive infection with immature parasites, sudden death could occur and lead to serious economic losses (Torgerson & Claxton, 1999; Rojo-Vázquez *et al.*, 2012).

Severe infection with immature paramphistomes can cause great economic losses due to reduced weight gain and decrease in milk production (Horak, 1971). Paramphistomosis has been reported to cause between

30 and 40% mortality in cattle and sheep. These deaths may result from anaemia, hypo-proteinaemia, profuse diarrhoea and marked emaciation (Soulsby, 1982), and a decrease in milk yield in dairy cattle (Spence *et al.*, 1996).

Economic losses from dicrocoeliasis are less apparent compared to those from other trematode infections and are difficult to quantify, due to concurrent infection with other gastrointestinal parasites. This may also be largely due to the asymptomatic nature of the disease (Otranto & Traversa, 2003). Clinical signs are mostly absent, but losses occur mainly from liver condemnation due to fibrosis of bile ducts and cirrhosis (Taylor *et al.*, 2007).

An abattoir survey to determine the prevalence and species of cattle schistosomiasis in Zambia revealed an overall prevalence of 51%, with 93% of the infected animals having up to 100 worm pairs in the mesenteric veins; *S. mattheei* was the predominant species (75%); while *S. leiperi* (12%) and *S. margrebowiei* (2%) were also reported (De Bont *et al.*, 1994). Schistosomiasis due to *S. bovis* has also been reported with a prevalence rate of 4.8% in Tanzania (Nzalawahe *et al.*, 2015). These parasites cause economic losses due to their long-term effect on growth and productivity, and also due to increased susceptibility to other parasitic and bacterial diseases (De Bont & Vercruysse, 1998). The clinical signs associated with the intestinal and hepatic form of disease in ruminants are haemorrhagic enteritis, haematuria, anaemia and loss of weight, with death within a few months of the disease. Nasal schistosomiasis, on the other hand, is a chronic disease that causes coryza and dyspnoea in infected animals (Aiello, 1998). Production losses reported due to schistosomiasis in cattle are attributed mainly to losses occurring in animals aged between 6 and 30 months, and are due to reduced weight gain, liver condemnation, poor future reproductive performance and death (McCauley *et al.*, 1984). Also, approximately 165 million cattle are likely to be infected with *Schistosoma* species worldwide (De Bont *et al.*, 1994).

Many trematode infections are also zoonotic, and thus a public health concern (Wolfe, 1966; Chen & Mott, 1990; Mas-Coma *et al.*, 2005). The World Health Organization has estimated that about 56 million people worldwide are infected by at least one zoonotic trematode species (Furst *et al.*, 2012) and up to 750 million people are at risk of infection (Keiser & Utzinger, 2009). It has been suggested that recent environmental changes associated with global warming, which favours the development of snail intermediate hosts, contribute to an increase in disease outbreaks (Mitchell, 2002).

Epidemiology of bovine fasciolosis in Nigeria

Tropical fasciolosis due to *F. gigantica* Cobbold 1855 has been reported in several parts of Africa (Schillhorn van Veen, 1980; Phiri *et al.*, 2005a; Abebe *et al.*, 2010; Nzalawahe *et al.*, 2014) and is likely to be a problem throughout the continent. The two species of *Fasciola* – *F. gigantica* and *F. hepatica* – and their respective snail intermediate hosts have both been reported in Africa. Fasciolosis is endemic in Nigeria and is of great economic importance (Ogunrinade & Ogunrinade, 1980). The disease has been reported with varying prevalence across

the country and is one of the most common causes of liver condemnation in abattoirs. The prevalence distribution of bovine fasciolosis is mapped (fig. 1) according to available data published between 1980 and 2016 (Schillhorn van Veen, 1980; Nwosu & Srivastava, 1993; Ngwu *et al.*, 2004; Opara, 2005; Ekwunife & Eneanya, 2006; Adedokun *et al.*, 2008; Umar *et al.*, 2009; Ibironke & Fasina, 2010; Sugun *et al.*, 2010; Gboeloh, 2012; Omoleye, 2012; Odigie & Odigie, 2013; Abraham & Jude, 2014; Ardo *et al.*, 2014; Magaji *et al.*, 2014; Ngele & Ibe, 2014; Onyeabor, 2014; Yahaya & Tyav, 2014; Ejeh *et al.*, 2015; Elelu *et al.*, 2016). The high prevalence rate from liver condemnation in a 3-year period reported in Lagos State (Ibironke & Fasina, 2010) can be explained by the fact that this State is the commercial nerve centre of Nigeria, with a very large human population, and many cattle from all over the country are often slaughtered at the abattoir.

Cattle total losses from *Fasciola* infection alone in Nigeria have been estimated to cost £32.5 million (Fabiya & Adeleye, 1982). A survey of abattoirs in Nigeria showed that about 70% of organ condemnation, mainly of livers, was due to fasciolosis (Alonge & Fasanmi, 1979). A recent study reported up to 88.1% liver condemnation due to liver flukes alone in Lagos abattoir (Ibironke & Fasina, 2010). These studies are in agreement with a 3-year abattoir study in south-western Nigeria in which losses from liver condemnation were estimated at US\$134,000 (Ibironke & Fasina, 2010). These data translate into huge economic losses from cumulative condemnation across the country.

Acute fasciolosis due to the migration of juvenile flukes is rare in cattle; however, a few cases have been reported in Nigeria. For example, on a dairy farm in north-central Nigeria, a pregnant cow was reported to have died during parturition, where post-mortem examination revealed inflammatory lesions and fibrosis due to fasciolosis (Okaiyeto *et al.*, 2012). The chronic form of the disease is the most common in cattle in Nigeria (Ogunrinade & Adegoke, 1982) and it occurs when small numbers of flukes finally enter the bile duct and infection becomes patent. This results in a chronic wasting disease from slow acquisition of liver flukes for months or even years. The normal tissues of the liver are replaced by fibrous tissues, with hyperplastic cholangitis and macroscopically visible calcification of the bile ducts, giving rise to so-called 'pipe-stem liver' (Soulsby, 1982; Taylor *et al.*, 2007). Clinically, the condition is characterized by weight loss, anaemia and oedema at the jaw (bottle jaw), thoracic and lower abdominal regions (Radostits *et al.*, 2000; Mitchell, 2002; Taylor *et al.*, 2007; Rojo-Vázquez *et al.*, 2012).

Factors influencing outbreaks of bovine fasciolosis

Season of the year

There are varying reports on the seasonal infection rates of *F. gigantica* across Nigeria. The majority of studies reported high infection rates at the beginning of the dry season (Schillhorn van Veen *et al.*, 1980; Umar *et al.*, 2009), which is in agreement with studies in other parts of




Fig. 1. Geographical distribution of bovine fasciolosis across different states of Nigeria (1980–2016) from published prevalence data (%) based on abattoir records, liver and coprological examination. States with 0% had no available data at the time of review.

Africa, such as Zambia, where bovine fasciolosis was also reported, with higher fluke abundance reported in the post-rainy season (Phiri *et al.*, 2005b). Similar observations were made in Zimbabwe, where *Fasciola* faecal egg counts were reported to follow a seasonal variation, with an increase during the end of the dry season and highest liver condemnation reported during the rainy season (reviewed by Pfukenyi *et al.*, 2005). However, another study carried out in the southern part of Nigeria reported an all-year occurrence of infection (Gboeloh, 2012). This all-year occurrence in the south was attributed to the favourable climatic condition (warm and humid), which favours development of parasites and the increase in the density of snail intermediate hosts (Gboeloh, 2012). The climatic conditions in southern Nigeria could be likened to those of highland areas of Iringa, Tanzania (Nzalawahe *et al.*, 2015) and Zimbabwe (Pfukenyi *et al.*, 2006a), where wet/swampy grazing areas reportedly favour the availability and distribution of the snail intermediate host. In addition, a recent study in Tanzania has associated trematode infections, including *Fasciola*, with irrigation practices during the dry season that favour growth of intermediate snail hosts and development of trematode larval stages (Nzalawahe *et al.*, 2014).

Type of management system

The type of management system has been shown to influence significantly the prevalence of *Fasciola* infection (Keyyu *et al.*, 2005). A low prevalence rate was also reported to occur in cattle reared by a sedentary husbandry system in Lake Chad area (Jean-Richard *et al.*, 2014). While a high prevalence rate of 54.3% has been reported in cattle managed extensively in south-western Nigeria (Adediran

et al., 2014), this is not surprising, because the cattle come into contact with a snail-infected habitat during extensive communal grazing. In contrast, under intensive management systems, metacercaria-free water and herbage can be supplied to the cattle, thereby minimizing the likelihood of outbreaks of fasciolosis.

Age of cattle

In a recent study from Nigeria, *F. gigantica* infection was reported to be higher in adult cattle than weaners (Elelu *et al.*, 2016). This is in agreement with past studies in Africa with similar findings (Pfukenyi *et al.*, 2006a; Nzalawahe *et al.*, 2014). The difference in prevalence within age group was attributed to a longer length of exposure to the infection in adults compared to weaners (Pfukenyi *et al.*, 2005; Nzalawahe *et al.*, 2014). It was also concluded that adult cattle act as a constant source of *F. gigantica* infection for the more susceptible young animals (Pfukenyi *et al.*, 2005).

Distribution of the snail intermediate host

The intermediate snail host *Lymnaea (Radix) natalensis*, which is widespread throughout Africa, is also widely distributed in Nigeria, although its occurrence is restricted to permanent water bodies (Ndifon & Ukoli, 1989). *Lymnaea* is fairly common in regions with rainfall over 1000 mm (Schillhorn van Veen, 1980) and it tolerates relatively high temperatures (Njoku-Tony, 2011) consistent with tropical climates. In Uganda, *L. (Radix) natalensis*, the intermediate snail host of *F. gigantica*, is the vector commonly found abundantly at lower altitudes below 1800 m, while *Galba truncatula* (intermediate snail host of

F. hepatica) was found only at altitudes above 3000 m (Howell *et al.*, 2012). The presence of the two vectors indicates the possibility of the two *Fasciola* parasites in Uganda. Other countries in Africa have reported the presence of the two species of *Fasciola* (Yilma & Mesfin, 2000; Walker *et al.*, 2008; Sisay & Nibret, 2013). To date, the authors of this review are not aware of any report of *G. truncatula* in Nigeria, although specific studies to identify these species in areas of high altitude might be worthwhile.

In Nigeria, during the wet season, many of the developing *Lymnaea* snails are washed away in torrential streams after heavy rain and this may be important in the spread of *Fasciola* (Schillhorn van Veen, 1980). Snails were also reported to be more abundant during the beginning of the dry season, to reach a climax in the middle of the dry season, but to decrease towards the end of the dry season when most streams and pools dry up (Schillhorn van Veen, 1980). This is supported by a field study in south-western Nigeria showing that dry-season conditions favour snails, and this was said to be due to low turbidity, reduced currents and substantial growths of algae and macrophytes (Ndifon & Ukoli, 1989). In the sub-tropical and tropical countries with distinct wet and dry seasons, optimal development of fluke eggs to miracidia occurs at the start of the wet season and development within the snail is complete by the end of the rains (Taylor *et al.*, 2007). The dry season therefore coincides with the snail shedding of cercaria and more animals grazing closer to streams and ponds, thereby predisposing them to infection. Herdsmen migrate in search of water and grazing during the dry season, and thousands of cattle often converge on the few ponds that fail to dry up (Ikeme & Obioha, 1973).

Diagnosis of bovine fasciolosis

The most common method of diagnosis is by faecal egg counts and pathological lesions in the liver during abattoir examination. Serological diagnostic methods to detect antibodies, such as indirect enzyme-linked immunosorbent assay (ELISA) (Damwesh & Ardo, 2013; Aliyu *et al.*, 2014) and direct ELISA (Fagbemi *et al.*, 1997), have been carried out in Nigeria. Testing for precipitating antibodies using the Agar Gel Precipitation Test (AGPT) is another method of diagnosis (Adedokun *et al.*, 2008). Comparatively, this method has been shown to detect more positive cases than faecal and bile egg counts (Adedokun *et al.*, 2008). In Kwara State, north-central Nigeria, a survey of cattle for fasciolosis revealed higher prevalence rates from faecal analysis compared to a previous abattoir study that utilized liver examination. This further showed the lack of sensitivity of the abattoir method of diagnosis, as positive samples are likely to be lost (Elelu *et al.*, 2016).

Epidemiology of bovine paramphistomosis in Nigeria

Rumen flukes are paramphistome parasites commonly seen in the abattoir in Nigeria (Bunza *et al.*, 2008). The snail vector of the disease (*Lymnaea*, *Planorbis* and

Bulinus) has also been reported in Nigeria (Ndifon & Ukoli, 1989; Brown & Kristensen, 1993). The species reported to occur in Nigerian domestic livestock include *P. microbothrium*, *Carmynerius gregarious*, *Carmynerius spatio-sus*, *Cotylophorum cotylophorum* (Schillhorn van Veen *et al.*, 1975) and *P. cervi* (Bogatko, 1975). Other species recovered from Nigerian cattle are *Ceylonocotyle dicranocoelium*, *Bothriophoron bothriophoron*, *Calicophoron calicophorum* and *Calicophoron microbothrioides* (Dube *et al.*, 2013).

Prevalence rates of 2.2% (Edosomwan & Shoyemi, 2012); 18.8% (Nwigwe *et al.*, 2013) and 16.1% (Elelu *et al.*, 2016) have been reported for paramphistomes in past studies in Nigeria. A higher prevalence rate of 41.67% for *P. cervi* was reported in another abattoir study in northern Nigeria (Nnabuife *et al.*, 2013). Similarly, also in northern Nigeria, a high paramphistome prevalence rate of up to 56.0% in slaughtered cattle has been reported in a study area around a marshy river valley. The grazing land associated with this river valley provides suitable breeding sites for snail intermediate hosts of the parasites (Bunza *et al.*, 2008). The convergence of cattle in common grazing land during the dry season, as well as irrigation practices, have been implicated in the high prevalence of paramphistomosis (Nzalawahe *et al.*, 2014). In a cross-sectional study in Ethiopia, 51.8% of slaughtered cattle were positive for paramphistomosis, with peak prevalence rates observed during October to November (Ayalew *et al.*, 2016). Although adult cattle were more likely to be infected with paramphistomes because they are more likely to be infected when taken out for grazing, several studies have shown that there is no statistically significant difference in paramphistome infection between age groups of cattle (Titi *et al.*, 2010; Khedri *et al.*, 2015; Ayalew *et al.*, 2016).

Epidemiology of bovine dicrocoeliasis in Nigeria

There are records of *Dicrocoelium hospes* infection in cattle in Nigeria (Nwosu & Srivastava, 1993; Ulayi *et al.*, 2007), and snail species of *Limnicolaria flammea* have been shown experimentally to be a suitable intermediate host of the disease (Fashuyi & Adeoye, 1986). The majority of prevalence data are based on abattoir surveys (FAO, 1992). Studies carried out in Zaria abattoir, northern Nigeria recorded high prevalence rates of 56.0% (Schillhorn van Veen *et al.*, 1980) and 35.4% (Ulayi *et al.*, 2007) for *D. hospes* from cattle. A low prevalence rate of 18.3% was reported in Borno State (Nwosu & Srivastava, 1993), and 22.33% in Plateau State in the bile duct of cattle (Omowaye *et al.*, 2012). Seasonal infection rates have also been reported for *D. hospes* in Nigeria, with the highest rate occurring during and directly after the rainy season (Schillhorn van Veen *et al.*, 1980).

Epidemiology of bovine schistosomiasis in Nigeria

Schistosoma curassoni and *S. bovis* have been reported in cattle in Nigeria (Ndifon *et al.*, 1988; Elelu *et al.*, 2016). The intermediate snail host (*Bulinus globosus*) of *Schistosoma* species is present in Nigeria (Ndifon & Ukoli, 1989) and

has been experimentally infected with *S. bovis* miracidia originating from a Nigerian cow (Ndifon *et al.*, 1988). The transmission of *S. matthei* has been reported, with occurrence throughout the year and with high prevalence during the wet months (Pfukenyi *et al.*, 2005). The prevalence rates reported in Nigeria were 7.8% for *S. bovis* and 2.2% for *S. curassoni*, based on examination of rectal scrapings; however, a higher prevalence of 31.1% (including both species of *Schistosoma*) was observed in 502 slaughtered cattle by examination of mesenteric and rectal veins (Ndifon *et al.*, 1988). In another study that utilized a more sensitive ELISA technique, a higher prevalence rate (16.2%) was recorded in cattle of the Kuri breed compared with other breeds (Hambali *et al.*, 2016). This is not surprising as this breed of cattle, with their characteristic bulbous horns, is adapted to swimming; they are therefore more likely to come in contact with *Schistosoma*-infected water. Schistosomiasis is transmitted via active skin penetration during contact with water. Other studies carried out in parts of north-western Ethiopia, located close to swampy areas, reported higher prevalence rates of up to 24% in cattle (Lulie & Guadu, 2014). In addition, a significantly higher schistosomiasis prevalence was observed in the wet season compared to the dry season in cattle in Zimbabwe (Pfukenyi *et al.*, 2006b). A higher prevalence rate (25.2%) has also been reported in cattle managed extensively compared to those under a semi-intensive management system (15.38%) (Lulie & Guadu, 2014).

Molecular identification of trematode species

Several polymerase chain reaction (PCR)-based techniques have been used to identify digenetic trematodes (Lotfy *et al.*, 2008, 2010). Molecular identification has an important contribution to make in many areas where there are problems associated with definitive speciation. These areas include instances of species overlap as seen in *Fasciola* (Kendall, 1965); the existence of hybrids between different genotypes, the Japanese triploid forms for example (Itagaki & Tsutsumi, 1998); the identification of sexually immature paramphistome species (Horak, 1971); and difficulty in speciation based on morphology of parasite eggs (Itagaki *et al.*, 2003). Other reasons for molecular identification are the close relatedness between taxa, as seen between *S. haematobium* and *S. bovis* (human and animal parasites), as well as difficulty in cercarial identification (Webster *et al.*, 2010).

Molecular studies by analysis of the first (ITS1) and second (ITS2) internally transcribed spacer of the ribosomal rDNA and the mitochondrial cytochrome *c* oxidase I (COI) gene have been carried out for *Fasciola* (Mas-Coma *et al.*, 2005; Ali *et al.*, 2008; Amer *et al.*, 2011; Amor *et al.*, 2011), paramphistome (Lotfy *et al.*, 2010) and *Dicrocoelium* (Otranto *et al.*, 2007). ITS-1 and ITS-2 are highly conserved and useful for differentiating closely related taxa that have diverged relatively recently (<50 million years ago) (Mas-Coma & Bargues, 2009). The mitochondrial NADH dehydrogenase subunit 1 and COI were used recently to characterize *F. gigantica* from Nigeria (Ichikawa-Seki *et al.*, 2017). However, further molecular studies from different parts of Nigeria are

recommended in order to understand these important pathogens.

Other genetic methods, such as the use of restriction fragment length polymorphism (PCR-RFLP) of ribosomal or mitochondrial genes using common restriction enzymes (such as *Ava*II and *Dra*II), have been used to analyse the whole of mitochondrial DNA to distinguish *Fasciola* species (Marcilla *et al.*, 2002), to identify species of paramphistomes (Itagaki *et al.*, 2003), to study genetic variability of *Dicrocoelium* species (Sandoval *et al.*, 1999), and also to distinguish between *S. haematobium* and *S. bovis* (Barber *et al.*, 2000; Webster *et al.*, 2010).

A PCR technique has also been used to detect the *F. gigantica* infection status of snail intermediate hosts (Velusamy *et al.*, 2004; Kaset *et al.*, 2010). A PCR technique used to amplify specific fragments of mitochondrial DNA in faecal samples of sheep has shown promise in the early detection of *F. hepatica* infection (Martínez-Pérez *et al.*, 2012). Molecular identification of paramphistome species from Asia and elsewhere in Africa has been carried out (Lotfy *et al.*, 2010); however, at time of writing this review, there are no records of such studies having been carried out in Nigeria. However, PCR to amplify species-specific genes has been used in differentiating human species of *Schistosoma* in Nigeria (Akinwale *et al.*, 2014) and also to identify *S. bovis* in Kenya (Kamanja *et al.*, 2011).

Control strategies for trematode infections

The principles behind the control of trematode infections are similar for all species (FAO, 1993). Conventional methods for control include strategic anthelmintic treatment to reduce environmental contamination (FAO, 1993), snail eradication by use of molluscicides and improved drainage systems to adversely influence the snail habitats (Armour, 1975; De Bont & Vercruysse, 1997). The current control of *F. hepatica* in cattle in temperate regions where the disease is prevalent is often based on strategically timed flukicide treatment, which is determined by studying the seasonal transmission dynamics in numerous locations throughout the world (Kaplan, 2001). The tropical trematode infection caused by *F. gigantica* may be less amenable to this approach because the intermediate hosts *L. (Radix) natalensis* are true water snails (Sewell, 1966) and cattle can come into contact with infected snails while grazing around water bodies throughout the year. Control of *Dicrocoelium* infection is more difficult due to the complex life cycle, involving terrestrial intermediate hosts. Control is therefore based on chemotherapy and husbandry practices (avoiding grazing during early morning and late evening to avoid infective ants) (Otranto & Traversa, 2002).

Strategic chemotherapy

Strategic anthelmintic treatment based on epidemiological and meteorological data is important for the control of flukes (FAO, 1993). There are several drugs for the treatment of fasciolosis. These include: halogenated phenol (niclofolan, bithionol, hexachlorophene and nitroxylin), salicylanides (rafoxanide, oxcyclozanide and

closantel), benzimidazoles (triclabendazole and albendazole), sulphonamides (clorsulon) and phenoxyalkanes (diamphenethides) (Fairweather & Boray, 1999a). Triclabendazole has been the preferred drug for treating liver flukes since 1983, due to its high efficacy against early immature, immature and adult flukes (Boray *et al.*, 1983). These drugs differ in their effectiveness against adults and immature flukes. Strategic antihelminthic use at 12- to 13-week intervals is effective against both mature and immature flukes and reduces the intensity of infection over time. Suggestions for strategic antihelminthic treatments have been suggested: in the tropics, with fasciolosis outbreaks all year round, treatment up to four times per year is recommended (Torgerson & Claxton, 1999). In Nigeria, some parts have reported seasonal trends in fasciolosis, while some southern parts of the country have reported an all-year-round occurrence (Gboeloh, 2012). Some authorities recommend that cattle should be dewormed regularly (Aliyu *et al.*, 2014), while others recommend treatment upon onset of clinical fasciolosis. Damwesh & Ardo (2015) proposed 2–3 annual treatments: at the start of the rainy season, mid rainy season and at the start of the dry season.

The anthelmintic drugs currently in use in Nigeria include albendazole, nitroxylnil, closulon and levamisole. However, a search in the available literature revealed that the preferred drug of choice against *Fasciola* (triclabendazole) is currently not available for use in Nigeria.

Neither triclabendazole nor niclofolan have been found to be effective against *Dicrocoelium* and paramphistome species (Güralp & Tinar, 1984). The drugs of choice for paramphistome infection are resorantel, oxcyclozanide and the combination of bithional and levamisole (Aiello, 1998). A combination of oxcyclozanide and levamisole is most effective against paramphistomosis in cattle, given in two treatments 3 days apart (Rolfe & Boray, 1987). Strategic treatment during the dry season may reduce contamination of the snails' habitat in the following rainy season (Rolfe *et al.*, 1991).

Benzimidazoles (except triclabendazole) and probenzimidazoles (thiophanate, netobimin) at higher doses are effective against *Dicrocoelium* species (Otranto & Traversa, 2002). Praziquantel is highly effective against all bovine visceral schistosomiasis (De Bont & Vercruysse, 1997) but is advised only in severe outbreaks due to the risk of portal occlusion from heavy worm burdens (McCully & Kruger, 1969). A combination of praziquantel and artemether is also effective against mature and immature schistosomes (Pfunkenyi *et al.*, 2005).

Anthelmintic drug resistance

Resistance of parasites to anthelmintic drugs is a growing global problem (Wanyangu *et al.*, 1996). Anthelmintic drug resistance is a real problem in the UK (Gordon *et al.*, 2012) and Australia (Brockwell *et al.*, 2014), and has also been reported in humans in The Netherlands (Winkelhagen *et al.*, 2012). Anthelmintic resistance has been demonstrated using faecal egg count reduction (Coles *et al.*, 1992), copro-antigen reduction tests (Brockwell *et al.*, 2014) and egg-hatch assay (Fairweather *et al.*, 2012). Demonstration of resistance to triclabendazole (also the drug of choice in human fasciolosis) against

immature and adult *Fasciola* indicates serious potential problems in controlling fasciolosis in the future (Fairweather & Boray, 1999b). Burdens of 20–34 drug-resistant flukes were observed in cattle following treatment with triclabendazole in Australia (Brockwell *et al.*, 2014). However, studies have shown that the anthelmintic drugs nitroxylnil and oxcyclozanide were able to kill 100% and 99.6% of adult triclabendazole-resistant flukes, respectively. Also, albendazole caused up to 95% reduction in triclabendazole-resistant fluke egg counts, while clorsulon showed a 73.2% reduction (Coles & Stafford, 2001). This finding is consistent with those carried out recently, showing that the anthelmintic drugs clorsulon and oxcyclozanide were effective in removing adult triclabendazole-resistant flukes (Elliott *et al.*, 2015).

There are also reports of some isolates of *F. hepatica* being resistant to albendazole in sheep (Novobilsky *et al.*, 2012; Sanabria *et al.*, 2013). A reduced efficacy of albendazole and oxcyclozanide against *F. gigantica* in naturally infected cattle has also been reported in Tanzania (Keyyu *et al.*, 2008). This could probably be the case with *F. gigantica* in Nigeria, hence further validation tests may be needed. There are currently few, if any, data on anthelmintic drug resistance in Nigerian cattle. However, a recent study in small ruminants suggests low resistance to ivermectin and levamisole with susceptibility to albendazole (Adediran & Uwalaka, 2015).

The incomplete elimination of infection leads to subclinical fasciolosis with continuous contamination of pasture, especially in sheep (Ollerenshaw, 1971). Use of a combination of drugs – synergistic drug usage (Fairweather & Boray, 1999b) – and treatment based on degree of infection (Malone & Craig, 1990) may be practical treatment strategies.

Human trematodiasis in Nigeria

Human fasciolosis has been reported from 51 countries over the past 25 years in the continents of Africa, America, Asia, Europe and Oceania (Mas-Coma *et al.*, 1999). Fasciolosis is among the most neglected tropical diseases (Mas-Coma *et al.*, 2009), with an estimated 180 million humans at risk of infection by *Fasciola* species worldwide (WHO, 1995) and with as many as 2.4–17 million humans infected (Hopkins, 1992; Toledo *et al.*, 2012). Human infection with fasciolosis has been reported to occur from eating uncooked watercress derived from endemic areas where infected cattle range freely, and also probably from contaminated water (Stemmermann, 1953; Toledo *et al.*, 2012). Human fasciolosis is determined by the presence of the intermediate snail host, domestic herbivorous animals, climatic conditions and the dietary habits of man (Chen & Mott, 1990). Mas-Coma *et al.* (1999) have reviewed the prevalence of *Fasciola* species reported in various parts of the world.

The main methods of diagnosing human trematodiasis are direct parasitological detection of fluke eggs in stools (Mas-Coma *et al.*, 2006) and other biofluids (duodenal and biliary aspirates), and immune-diagnosis (indirect diagnosis). Other non-invasive diagnostic techniques, such as radiology, ultrasound, computed tomography and magnetic resonance imaging, can also be used

(Esteban *et al.*, 1998; Mas-Coma *et al.*, 2005; Keiser & Utzinger, 2009). In a recent review, stool and blood techniques have been reported to be improved for diagnosis of human fasciolosis and during surveys. However, the author also identified difficulties of diagnosing fascioliasis in humans due to different infection phases and parasite migration capacities, clinical heterogeneity, immunological complexity, different epidemiological situations and transmission patterns (Mas-Coma *et al.*, 2014).

Human fasciolosis involving both species of *Fasciola* has been reported in Africa (Mas-Coma *et al.*, 1999). There are hardly any studies targeted at detecting the zoonotic effect of fasciolosis in Nigeria. There is a report of the positive detection of *Fasciola* species in HIV-infected human subjects in Nigeria, although at a low prevalence rate of 1% (Abaver *et al.*, 2012). This suggests that human infection may be significant, although further research on risk factors as well as possible routes of transmission needs to be carried out.

Human cases of microcoeliasis are rare and often occur by accidental ingestion of infected ants on unwashed vegetables, or by drinking contaminated water (Haridy *et al.*, 2003). Human cases of microcoeliasis have been reported in Nigeria (Roche, 1948; Samaila *et al.*, 2009), Czechoslovakia (Ondriska *et al.*, 1989), Egypt (Massoud *et al.*, 2003), Turkey (Cengiz *et al.*, 2010) and Kyrgyzstan (Jeandron *et al.*, 2011). Spurious infection with *D. hospes* has also been reported in Ghana due to accidental ingestion of infected animal liver (Wolfe, 1966). The subclinical form of the disease, which is most common, is characterized by cholangitis and adenomatous proliferation of the bile duct (Cabeza-Barrera *et al.*, 2011). Other signs are constipation, diarrhoea, vomiting and abdominal pain (Ondriska *et al.*, 1989). Human microcoeliasis has been reported in a 7-year-old Nigerian child of a nomadic pastoralist, showing clinical signs of fever, jaundice and anterior subcutaneous abdominal mass (Samaila *et al.*, 2009).

The amphistome species reported in man is *Gastrodiscoides hominis*, which is widely distributed in Asia. It is located in the caecum and colon, with pigs being the main animal reservoir (Dutt & Srivastava, 1972) and has been reported in Nigeria in a malnourished child (Dada-Adegbola *et al.*, 2004).

Although *S. bovis* primarily infects ruminants, it has been isolated in humans in various parts of Africa (Raper, 1951; Chunge *et al.*, 1986; Kinoti & Mumo, 1988; Santoro, 1988). Moreover, the mesenteries of cattle, which may be infected with *S. bovis*, are usually sold at the local markets in Africa as edible tripe, thus making it a public health risk (Kamanja *et al.*, 2011). Hybridization between *S. bovis* and *S. haematobium* (cattle and human infection) has been reported in Senegal, with an impending risk of the emergence of a new disease (Huyse *et al.*, 2009).

Several drugs for the effective treatment of human fasciolosis are available. These include dihydroemetine-emetine derivative, bithionol, praziquantel and triclabendazole (effective against both acute and chronic fasciolosis). These are currently the drugs of choice for human fasciolosis caused by both *F. hepatica* and *F. gigantica* (Esteban *et al.*, 1998; Savioli *et al.*, 1999). Praziquantel has also been used in the treatment of human microcoeliasis (Massoud *et al.*, 2003). Mebendazole was effective in treatment of the human amphistomosis (Dada-Adegbola *et al.*, 2004).

The prevention of human trematode infections may be achieved by strict control of water cress and other metacercariae-carrying aquatic plants for human consumption, especially in endemic zones (Mas-Coma *et al.*, 2005). Drinking water must be boiled or purified. Integrated control approaches and inter-sectoral collaboration between public health and veterinary medicine have also been suggested for control (Keiser & Utzinger, 2009).

Conclusions and recommendations

Trematode infections are a significant limiting factor in livestock production; therefore, the development of sustainable strategies for their control is a priority. In order to develop sustainable control, gaps in knowledge must be identified to guide research projects. The outcome of such research would enhance proper allocation of funds for disease control and policy formulation. In addition, it is also important to carry out detailed study on the overall economic losses due to trematode infections, in order to design and implement appropriate systematic disease prevention and control methods.

The prevalence rates of trematodes reported across the various abattoirs in Nigeria are an indication that trematode infection poses a widespread risk to livestock and, possibly, humans. However, it is worthy of note that neither oxcyclozanide nor triclabendazole, which are the two drugs of choice for mixed and human *Fasciola* infection, respectively, is currently available for use in Nigeria. Availability of these two drugs could be helpful in reducing disease occurrence. These drugs could be made available by formulating favourable policies to ensure that drug retailers have basic training in identifying and stocking the appropriate drugs of choice and informing livestock farmers on the rationale for their use (an example is information on effectiveness of triclabendazole against both mature and immature flukes). The gap in technical knowledge needed by drug retailers for dispensing drugs has been identified in a previous study (Bett *et al.*, 2004) and veterinarians have a huge responsibility for advising/training these retailers on the appropriate drugs to stock, based on the diseases prevalent to different localities.

There are hardly any studies targeted at detecting the zoonotic effect of fasciolosis infections in Nigeria. Although a few studies exist on other trematodes in Nigeria, these are often case reports. Further research on burden and risk factors, as well as possible routes of transmission of trematode infection to humans, needs to be carried out.

Molecular studies on trematodes in different species of livestock, including those targeted at understanding anthelmintic drug resistance, should also be carried out across Nigeria. These studies would be useful in providing more data on the diversity of these important trematodes.

Financial support

The research is funded by a PhD studentship from the Nigerian Tertiary Education Trust Fund, the Leverhulme–Royal Society African Award Scheme and the University of Bristol, UK.

Conflict of interest

None.

References

- Abaver, D.T., Nwobegahay, J.M., Goon, D.T., Iweriebor, B.C. & Khoza, L.B. (2012) Enteric parasitic infections in HIV-infected patients with low CD4 counts in Toto, Nigeria. *Pakistan Journal of Medical Sciences* **28**, 630–633.
- Abebe, R., Abunna, F., Berhane, M., Mekuria, S., Megersa, B. & Regassa, A. (2010) Fasciolosis: prevalence, financial losses due to liver condemnation and evaluation of a simple sedimentation diagnostic technique in cattle slaughtered at Hawassa Municipal abattoir, Southern Ethiopia. *Ethiopia Veterinary Journal* **14**, 39–51.
- Abraham, J.T. & Jude, I.B. (2014) Fascioliasis in cattle and goat slaughtered at Calabar Abattoirs. *Journal of Biology, Agriculture and Healthcare* **4**, 34–40.
- Abrous, M., Rondelaud, D. & Dreyfuss, G. (1996) *Paramphistomum daubneyi*: the development of redial generations in the snail *Lymnaea truncatula*. *Parasitology Research* **83**, 64–69.
- Adediran, O.A. & Uwalaka, E.C. (2015) Effectiveness evaluation of levamisole, albendazole, ivermectin, and *Vernonia amygdalina* in West African dwarf goats. *Journal of Parasitology Research*. doi: <http://dx.doi.org/10.1155/2015/706824>.
- Adediran, O.A., Adebisi, A.I. & Uwalaka, E.C. (2014) Prevalence of *Fasciola* species in ruminants under extensive management system in Ibadan southwestern Nigeria. *African Journal of Medicine and Medical Sciences* **43** (Suppl.), 137–141.
- Adedokun, O.A., Ayinmode, A.B. & Fagbemi, B.O. (2008) A comparative study of three methods for detecting *Fasciola* infections in Nigerian cattle. *Veterinarski Arhiv* **78**, 411–416.
- Agatsuma, T., Arakawa, Y., Iwagami, M., Honzako, Y., Cahyaningsih, U., Kang, S.Y. & Hong, S.J. (2000) Molecular evidence of natural hybridization between *Fasciola hepatica* and *F. gigantica*. *Parasitology International* **49**, 231–238.
- Aiello, S.E. (1998) *The Merck veterinary manual*. Whitehouse Station, New Jersey, USA, Wiley.
- Akinwale, O.P., Hock, T.T., Chia-Kwung, F., Zheng, Q., Haimo, S., Ezech, C. & Gyang, P.V. (2014) Differentiating *Schistosoma haematobium* from *Schistosoma grebowiei* and other closely related schistosomes by polymerase chain reaction amplification of a species specific mitochondrial gene. *Tropical Parasitology* **4**, 38.
- Ali, H., Ai, L., Song, H.Q., Ali, S., Lin, R.Q., Seyni, B., Issa, G. & Zhu, X.Q. (2008) Genetic characterisation of *Fasciola* samples from different host species and geographical localities revealed the existence of *F. hepatica* and *F. gigantica* in Niger. *Parasitology Research* **102**, 1021–1024.
- Aliyu, A.A., Ajogi, I.A., Ajanusi, O.J. & Reuben, R.C. (2014) Epidemiological studies of *Fasciola gigantica* in cattle in Zaria, Nigeria using coprology and serology. *Journal of Public Health* **6**, 85–91.
- Alonge, D.O. & Fasanmi, E.F. (1979) A survey of abattoir data in Northern Nigeria [cattle]. *Tropical Animal Health and Production* **11**, 57–62.
- Alvarez Rojas, C.A., Jex, A.R., Gasser, R.B. & Scheerlinck, J.-P.Y. (2014) Techniques for the diagnosis of *Fasciola* infections in animals: room for improvement. *Advances in Parasitology* **86**, 65–107.
- Amer, S., Dar, Y., Ichikawa, M., Fukuda, Y., Tada, C., Itagaki, T. & Nakai, Y. (2011) Identification of *Fasciola* species isolated from Egypt based on sequence analysis of genomic (ITS1 and ITS2) and mitochondrial (ND1 and COI) gene markers. *Parasitology International* **60**, 5–12.
- Amor, N., Farjallah, S., Salem, M., Lamine, D.M., Merella, P., Said, K. & Ben Slimane, B. (2011) Molecular characterization of *Fasciola gigantica* from Mauritania based on mitochondrial and nuclear ribosomal DNA sequences. *Experimental Parasitology* **129**, 127–136.
- Ardo, M.B., Aliyara, Y.H. & Lawal, H. (2014) Prevalence of bovine fasciolosis in major abattoirs of Adamawa State, Nigeria. *Bayero Journal of Pure and Applied Sciences* **6**, 12–16.
- Aregheore, E.M. (Ed.) (2009) *Country forage/forest resource profile Nigeria*. pp. 1–42. Rome, Food and Agriculture Organization of United Nations.
- Armour, J. (1975) The epidemiology and control of bovine fascioliasis. *Veterinary Record* **96**, 198–201.
- Armstrong, A. (1982) Economic impact of fascioliasis. *VIIIth World Congress on diseases of cattle*. pp. 1113–1117. Netherlands, World Association for Buiatrics.
- Ashrafi, K., Valero, M.A., Panova, M., Perigo, M.V., Massoud, J. & Mas-Coma, S. (2006) Phenotypic analysis of adults of *Fasciola hepatica*, *Fasciola gigantica* and intermediate forms from the endemic region of Gilan, Iran. *Parasitology International* **55**, 249–260.
- Ayalew, G., Tilahun, A., Aylate, A., Teshale, A. & Getachew, A. (2016) A study on prevalence of paramphistomum in cattle slaughtered in Gondar Elfora Abattoir, Ethiopia. *Journal of Veterinary Medicine and Animal Health* **8**, 107–111.
- Babalola, D.A. & Schillhorn van Veen, T.W. (1976). Incidence of fascioliasis in cattle slaughtered in Bauchi (Nigeria). *Tropical Animal Health and Production* **8**, 243–247.
- Barber, K.E., Mkoji, G.M. & Loker, E.S. (2000) PCR-RFLP analysis of the ITS2 region to identify *Schistosoma haematobium* and *S. bovis* from Kenya. *American Journal of Tropical Medicine and Hygiene* **62**, 434–440.
- Bett, B., Machila, N., Gathura, P.B., Mcdermott, J.J. & Eisler, M.C. (2004) Characterisation of shops selling veterinary medicines in a tsetse-infested area of Kenya. *Preventive Veterinary Medicine* **63**, 29–38.
- Bogatko, W. (1975) Mass mixed infection with *Paramphistomum cervi* and *Fasciola gigantica* in cattle in northern Nigeria. *Medycyna Weterynaryjna* **31**, 469–470.
- Boray, J.C. (1969) Experimental fasciolosis in Australia. *Advances in Parasitology* **7**, 95–210.
- Boray, J.C. (1985) Flukes in domestic animals. pp. 179–218 in Gaafar, S.M., Howard, W.E. & Marsh, R.E. (Eds) *Parasites, pests and predators*. New York, Elsevier.

- Boray, J., Crowfoot, P., Strong, M., Allison, J., Schellenbaum, M., Von Orelli, M. & Sarasin, G. (1983) Treatment of immature and mature *Fasciola hepatica* infections in sheep with triclabendazole. *Veterinary Record* **113**, 315–317.
- Brockwell, Y.M., Elliott, T.P., Anderson, G.R., Stanton, R., Spithill, T.W. & Sangster, N.C. (2014) Confirmation of *Fasciola hepatica* resistant to triclabendazole in naturally infected Australian beef and dairy cattle. *International Journal for Parasitology: Drugs and Drug Resistance* **4**, 48–54.
- Brown, D.S. & Kristensen, T.K. (1993) *A field guide to African fresh water snails: West African species*. Denmark, Danish Bilharzias Laboratory.
- Bunza, M.D.A., Ahmad, A. & Fana, S.A. (2008) Prevalence and fluke burden of paramphistomiasis in ruminants slaughtered at Sokoto Central Abattoir, Sokoto. *Nigerian Journal of Basic and Applied Sciences* **16**, 287–292.
- Cabeza-Barrera, I., Cabezas-Fernández, T., Salas Coronas, J., Vázquez Villegas, J. & Cobo, F. (2011) *Dicrocoelium dendriticum*: an emerging spurious infection in a geographic area with a high level of immigration. *Annals of Tropical Medicine and Parasitology* **105**, 403–406.
- Castro-Trejo, L., García-Vasquez, Z. & Casildo-Nieto, J. (1990) The susceptibility of lymnaeid snails to *Paramphistomum cervi* infections in Mexico. *Veterinary Parasitology* **35**, 157–161.
- CBN (1999) *Annual Report*. Central Bank of Nigeria.
- Cengiz, Z.T., Yilmaz, H., Dülger, A.C. & Çiçek, M. (2010) Human infection with *Dicrocoelium dendriticum* in Turkey. *Annals of Saudi Medicine* **30**, 159–161.
- Chen, M.G. & Mott, K.E. (1990) Progress in assessment of morbidity due to *Fasciola hepatica* infection: a review of recent literature. *Tropical Diseases Bulletin* **87**, 1–38.
- Chunge, R., Katsivo, M., Kok, P., Wamivea, M. & Kinoti, S. (1986) *Schistosoma bovis* in human stools in Kenya. *Transactions of the Royal Society of Tropical Medicine and Hygiene* **80**, 849.
- Coles, G.C. & Stafford, K.A. (2001) Activity of oxyclozanide, nitroxylnil, clorsulon and albendazole against adult triclabendazole resistant *Fasciola hepatica*. *Veterinary Record* **148**, 723–724.
- Coles, G.C., Bauer, C., Borgsteede, F.H.M., Geerts, S., Klei, T.R., Taylor, M.A. & Waller, P.J. (1992) World Association for the Advancement of Veterinary Parasitology (W.A.A.V.P.) methods for the detection of anthelmintic resistance in nematodes of veterinary importance. *Veterinary Parasitology* **44**, 35–44.
- Dada-Adegbola, H.O., Falade, C.O., Oluwatoba, O.A. & Abiodun, O.O. (2004) *Gastrodiscoides hominis* infection in a Nigerian – case report. *West African Journal of Medicine* **23**, 185–186.
- Damwesh, S.D. & Ardo, M.B. (2013) Detection of *Fasciola gigantica* antibodies using Pourquier ELISA kit. *Sokoto Journal of Veterinary Sciences* **11**, 43–48.
- Damwesh, S.D. & Ardo, M.B. (2015) Best periods for deworming cattle against fasciolosis in Nigeria (a tropical sub-saharan country with dry and wet seasons). *Journal of Veterinary Science and Technology* **6**, 270.
- De Bont, J. & Vercruysse, J. (1997) The epidemiology and control of cattle schistosomiasis. *Parasitology Today* **13**, 255–262.
- De Bont, J. & Vercruysse, J. (1998) Schistosomiasis in cattle. *Advances in Parasitology* **41**, 285–364.
- De Bont, J., Vercruysse, J., Southgate, V.R., Rollinson, D. & Kaukas, A. (1994) Cattle schistosomiasis in Zambia. *Journal of Helminthology* **68**, 295–299.
- Dinnik, J.A. (1964) Intestinal paramphistomiasis and *P. microbothrium* Fischöder in Africa. *Bulletin of Epizootic Diseases of Africa* **12**, 439–454.
- Dipeolu, M.A., Dipeolu, A. & Eruvbetine, D. (2000) The prevalence of fascioliasis in south western Nigeria (1986–91). *International Journal of Animal Sciences* **15**, 151–152.
- Dube, S., Onyedineke, N.E. & Aisien, M.S.O. (2013) *Ceylonocotyle*, *Bothriophoron*, and *Calicophoron* species parasitic in some Nigerian cattle. *Advances in BioResearch* **4**, 38–43.
- Dutt, S.C. & Srivastava, H.D. (1972) The life history of *Gastrodiscoides hominis* (Lewis and McConnel, 1876) Leiper, 1913 – the amphistome parasite of man and pig. *Journal of Helminthology* **46**, 35–46.
- Edosomwan, E.U. & Shoyemi, O.O. (2012) Prevalence of gastrointestinal helminth parasites of cattle and goats slaughtered at abattoirs in Benin City, Nigeria. *African Scientist* **13**, 109–114.
- Ejeh, E.F., Paul, B.T., Lawan, F.A., Lawal, J.R., Ejeh, S.A. & Hambali, I.U. (2015) Seasonal prevalence of bovine fasciolosis and its direct economic losses (del) due to liver condemnation at Makurdi abattoirs, North central Nigeria. *Sokoto Journal of Veterinary Sciences* **13**, 42–48.
- Ekwunife, C.A. & Eneanya, C.I. (2006) *Fasciola gigantica* in Onitsha and environs. *Animal Research International* **3**, 448–450.
- Elelu, N., Ambali, A., Coles, G.C. & Eisler, M.C. (2016) Cross-sectional study of *Fasciola gigantica* and other trematode infections of cattle in Edu Local Government Area, Kwara State, North-central Nigeria. *Parasites & Vectors* **9**, 470.
- Elliott, T.P., Kelley, J.M., Rawlin, G. & Spithill, T.W. (2015) High prevalence of fasciolosis and evaluation of drug efficacy against *Fasciola hepatica* in dairy cattle in the Maffra and Bairnsdale districts of Gippsland, Victoria, Australia. *Veterinary Parasitology* **209**, 117–124.
- Esteban, J.G., Bargues, M.D. & Mas-Coma, S. (1998) Geographical distribution, diagnosis and treatment of human fasciolosis: a review. *Research and Reviews in Parasitology* **58**, 13–42.
- European Environment Agency (2004) *Impacts of Europe's changing climate. An indicator-based assessment*. EEA Report No. 2.
- Fabiya, J.P. (1987) Production losses and control of helminths in ruminants of tropical regions. *International Journal for Parasitology* **17**, 435–442.
- Fabiya, J.P. & Adeleye, G.A. (1982) Bovine fascioliasis on the Jos Plateau, Northern Nigeria with particular reference to economic importance. *Bulletin of Animal Health and Production in Africa* **30**, 41–43.
- Fagbemi, B.O., Aderibigbe, O.A. & Guobadia, E.E. (1997) The use of monoclonal antibody for the immunodiagnosis of *Fasciola gigantica* infection in cattle. *Veterinary Parasitology* **69**, 231–240.

- Fairweather, I. & Boray, J.C. (1999a) Fasciolicides: Efficacy, actions, resistance and its management. *The Veterinary Journal* **158**, 81–112.
- Fairweather, I. & Boray, J.C. (1999b) Mechanisms of fasciolicide action and drug resistance in *Fasciola hepatica*. pp. 225–276 in Dalton, J.P. (Ed.) *Fasciolosis*. Wallingford, CABI Publishing.
- Fairweather, I., McShane, D.D., Shaw, L., Ellison, S.E., O'Hagan, N.T., York, E.A., Trudgett, A. & Brennan, G.P. (2012) Development of an egg hatch assay for the diagnosis of triclabendazole resistance in *Fasciola hepatica*: proof of concept. *Veterinary Parasitology* **183**, 249–259.
- FAO (1992) *Distribution and impact of helminth diseases of livestock in developing countries*. Rome, Italy, FAO Document Repository.
- FAO (1993) The epidemiology of helminthes parasites. Available at <http://www.fao.org/wairdocs/ILRI/x5492E/x5492e04.htm-2.5%20trematodes> (accessed 20 April 2017).
- Fashuyi, S.A. & Adeoye, G.O. (1986) The possible snail intermediate hosts of *Dicrocoelium hospes* in Nigeria. *Acta Veterinaria Brno* **55**, 85–88.
- Furst, T., Keiser, J. & Utzinger, J. (2012) Global burden of human food-borne trematodiasis: a systematic review and meta-analysis. *Lancet Infectious Diseases* **12**, 210–221.
- Gboeloh, L.B. (2012) Seasonal prevalence of *Fasciola gigantica* in slaughtered cattle in major abattoirs in Port Harcourt. *Advances in Agriculture, Sciences and Engineering Research* **2**, 336–340.
- Gordon, D.K., Zadoks, R.N., Stevenson, H., Sargison, N. D. & Skuce, P.J. (2012) On farm evaluation of the coproantigen ELISA and coproantigen reduction test in Scottish sheep naturally infected with *Fasciola hepatica*. *Veterinary Parasitology* **187**, 436–444.
- Güralp, N. & Tinar, R. (1984) Trematodiasis in Turkey: comparative efficacy of triclabendazole and niclofolan against natural infections of *Fasciola hepatica* and *F. gigantica* in sheep. *Journal of Helminthology* **58**, 113–116.
- Hambali, I.U., Adamu, N.B., Ahmed, M.I., Bokko, P., Mbaya, A.W., Tijjani, A.O., Biu, A.A., Jesse, F.F.A. & Ambali, A.G. (2016) Sero-prevalence of *Schistosoma* species in cattle in Maiduguri Metropolis and Jere Local Government Areas of Borno State, Nigeria. *Journal of Advanced Veterinary and Animal Research* **3**, 56–61.
- Hammond, J.A. (1972) Infections with *Fasciola* spp in wildlife in Africa. *Tropical Animal Health and Production* **4**, 1–13.
- Haridy, F.M., Morsy, T.A., Ibrahim, B.B. & Abdel-Aziz, A. (2003) A preliminary study on dicrocoeliasis in Egypt, with a general review. *Journal Egypt Society for Parasitology* **33**, 85–96.
- Hillyer, G.V. (2005) *Fasciola* antigens as vaccines against fascioliasis and schistosomiasis. *Journal of Helminthology* **79**, 241–247.
- Hopkins, D.R. (1992) Homing in on helminths. *American Journal of Tropical Medicine and Hygiene* **46**, 626–634.
- Horak, I.G. (1971) Paramphistomiasis of domestic ruminants. *Advances in Parasitology* **9**, 33–72.
- Howell, A., Mugisha, L., Davies, J., Lacourse, E.J., Claridge, J., Williams, D.J.L., Kelly-Hope, L., Betson, M., Kabatereine, N.B. & Stothard, J.R. (2012) Bovine fasciolosis at increasing altitudes: parasitological and malacological sampling on the slopes of Mount Elgon, Uganda. *Parasites & Vectors* **5**, 196.
- Huyse, T., Webster, B.L., Geldof, S., Stothard, J.R., Diaw, O.T., Polman, K. & Rollinson, D. (2009) Bidirectional introgressive hybridization between a cattle and human schistosome species. *PLoS Pathogen* **5**, e1000571.
- Ibironke, A.A. & Fasina, F.O. (2010) Socio-economic implications of bovine liver rejection in a major abattoir in south-western Nigeria. *Revista de Ciencias Agrarias* **33**, 211–216.
- Ichikawa-Seki, M., Tokashiki, M., Opara, M.N., Iroh, G., Hayashi, K., Kumar, U.M. & Itagaki, T. (2017) Molecular characterization and phylogenetic analysis of *Fasciola gigantica* from Nigeria. *Parasitology International* **66**, 893–897.
- Ikeme, M.M. & Obioha, F. (1973) *Fasciola gigantica* infestations in trade cattle in Eastern Nigeria. *Bulletin of Epizootic Diseases of Africa* **21**, 259–264.
- Itagaki, T. & Tsutsumi, K. (1998) Triploid form of *Fasciola* in Japan: genetic relationships between *Fasciola hepatica* and *Fasciola gigantica* determined by ITS-2 sequence of nuclear rDNA. *International Journal of Parasitology* **28**, 777–781.
- Itagaki, T., Tsumagari, N., Tsutsumi, K.-I. & Chinone, S. (2003) Discrimination of three amphistome species by PCR-RFLP based on rDNA ITS2 markers. *Journal of Veterinary Medical Science* **65**, 931–933.
- Jeandron, A., Rinaldi, L., Abdylidaieva, G., Usubalieva, J., Steinmann, P., Cringoli, G. & Utzinger, J. (2011) Human infections with *Dicrocoelium dendriticum* in Kyrgyzstan: the tip of the iceberg? *Journal of Parasitology* **97**, 1170–1172.
- Jean-Richard, V., Crump, L., Abicho, A.A., Naré, N.B., Greter, H., Hattendorf, J., Schelling, E. & Zinsstag, J. (2014) Prevalence of *Fasciola gigantica* infection in slaughtered animals in southeastern Lake Chad area in relation to husbandry practices and seasonal water levels. *BMC Veterinary Research* **10**, 81.
- Kamanja, I.T.G., Githigia, S.M., Muchemi, G.M. & Mwandawiro, C. (2011) A survey of *Schistosoma bovis* in cattle in Kwale district, Kenya. *Bulletin of Animal Health and Production in Africa* **59**, 161–168.
- Kaplan, R.M. (2001) *Fasciola hepatica*: a review of the economic impact in cattle and consideration for control. *Veterinary Therapeutics* **2**, 40–50.
- Kaset, C., Eursitthichai, V., Vichasri-Grams, S., Viyanant, V. & Grams, R. (2010) Rapid identification of *Lymnaea* snails and their infection with *Fasciola gigantica* in Thailand. *Experimental Parasitology* **126**, 482–488.
- Keiser, J. & Utzinger, J. (2009) Food-borne trematodiasis. *Clinical Microbiology Reviews* **22**, 466–483.
- Kendall, S.B. (1965) The relationships between the species of *Fasciola* and their molluscan hosts. *Advances in Parasitology* **3**, 59–98.
- Keyyu, J.D., Monrad, J., Kyvsgaard, N.C. & Kassuku, A. A. (2005) Epidemiology of *Fasciola gigantica* and amphistomes in cattle on traditional, small scale dairy and large scale dairy farms in the southern highlands of Tanzania. *Tropical Animal Health Production* **37**, 303–314.

- Keyyu, J.D., Kassuku, A.A., Kyvsgaard, N.C. & Monrad, J. (2008) Comparative efficacy of anthelmintics against *Fasciola gigantica* and amphistomes in naturally infected cattle in Kilolo District, Tanzania. *Tanzania Veterinary Journal* **25**, 40–47.
- Khedri, J., Radfar, M.H., Borji, H. & Mirzaei, M. (2015) Prevalence and intensity of *Paramphistomum* spp. in cattle from South-Eastern Iran. *Iran Journal of Parasitology* **10**, 268–272.
- Kinoti, G.K. & Mumo, J.M. (1988) Spurious human infection with *Schistosoma bovis*. *Transactions of the Royal Society of Tropical Medicine and Hygiene* **82**, 589–590.
- Lotfy, W.M., Brant, S.V., Dejong, R.J., Le, T.H., Demiaszkiewicz, A., Rajapakse, R.P., Perera, V.B., Laursen, J.R. & Loker, E.S. (2008) Evolutionary origins, diversification, and biogeography of liver flukes (Digenea, Fasciolidae). *American Journal Tropical Medicine & Hygiene* **79**, 248–255.
- Lotfy, W.M., Brant, S.V., Ashmawy, K.I., Devkota, R., Mkoji, G.M. & Loker, E.S. (2010) A molecular approach for identification of paramphistomes from Africa and Asia. *Veterinary Parasitology* **174**, 234–240.
- Lulie, B. & Guadu, T. (2014) Bovine schistosomiasis: A threat in public health perspective in Bahir Dar town, Northwest Ethiopia. *Acta Parasitologica Globalis* **5**, 01–06.
- Magaji, A., Ibrahim, K., Salihu, M.D., Saulawa, M.A., Mohammed, A.A. & Musawa, A.I. (2014) Prevalence of fascioliasis in cattle slaughtered in Sokoto Metropolitan Abattoir, Sokoto, Nigeria. *Advances in Epidemiology* **2014**.
- Malone, J.B. & Craig, T.M. (1990) Cattle liver flukes, risk assessment and control. *Compendium on Continuing Education for Practising Veterinarians* **12**, 747–754.
- Marcilla, A., Bargues, M.D. & Mas-Coma, S. (2002) A PCR-RFLP assay for the distinction between *Fasciola hepatica* and *Fasciola gigantica*. *Molecular and Cellular Probes* **16**, 327–333.
- Martínez-Pérez, J.M., Robles-Pérez, D., Rojo-Vázquez, F.A. & Martínez-Valladares, M. (2012) Comparison of three different techniques to diagnose *Fasciola hepatica* infection in experimentally and naturally infected sheep. *Veterinary Parasitology* **190**, 80–86.
- Mas-Coma, S. & Bargues, M.D. (2009) Populations, hybrids and the systematic concepts of species and subspecies in Chagas disease triatomine vectors inferred from nuclear ribosomal and mitochondrial DNA. *Acta Tropica* **110**, 112–136.
- Mas-Coma, M.S., Esteban, J.G. & Bargues, M.D. (1999) Epidemiology of human fasciolosis: a review and proposed new classification. *Bulletin of the World Health Organization* **77**, 340–346.
- Mas-Coma, S., Bargues, M.D. & Valero, M.A. (2005) Fascioliasis and other plant-borne trematode zoonoses. *International Journal Parasitology* **35**, 1255–1278.
- Mas-Coma, S., Bargues, M.D. & Valero, M.A. (2006) Gastrodiscoidiasis, a plant-borne zoonotic disease caused by the intestinal amphistome fluke *Gastrodiscoides hominis* (Trematoda: Gastrodiscidae). *Revista Ibérica de Parasitología* **66**, 75–81.
- Mas-Coma, S., Valero, M.A. & Bargues, M.D. (2009) Fasciola, lymnaeids and human fascioliasis, with a global overview on disease transmission, epidemiology, evolutionary genetics, molecular epidemiology and control. *Advances in Parasitology* **69**, 41–146.
- Mas-Coma, S., Bargues, M.D. & Valero, M.A. (2014) Diagnosis of human fascioliasis by stool and blood techniques: Update for the present global scenario. *Parasitology* **141**, 1918–1946.
- Massoud, A., Morsy, T.A. & Haridy, F.M. (2003) Treatment of Egyptian microcoeliasis in man and animals with Mirazid. *Journal of the Egyptian Society of Parasitology* **33**, 437–442.
- McCauley, E.H., Majid, A.A. & Tayeb, A. (1984) Economic evaluation of the production impact of bovine schistosomiasis and vaccination in the Sudan. *Preventive Veterinary Medicine* **2**, 735–754.
- McCully, R.M. & Kruger, S.P. (1969) Observation on bilharziasis of domestic ruminants in South Africa. *Onderstepoort Journal of Veterinary Research* **36**, 129–162.
- Mitchell, G. (2002) Update on fasciolosis in cattle. *In Practice* **24**, 378–385.
- Ndifon, G.T. & Ukoli, F.M.A. (1989) Ecology of freshwater snails in south-western Nigeria, I: distribution and habitat preferences. *Hydrobiologia* **171**, 231–253.
- Ndifon, G.T., Betterton, C. & Rollinson, D. (1988) *Schistosoma curassoni* Brumpt, 1931 and *S. bovis* (Sonsino, 1876) in cattle in northern Nigeria. *Journal of Helminthology* **62**, 33–34.
- Ngele, K.K. & Ibe, E. (2014) Prevalence of fasciolosis in cattle slaughtered at Eke market abattoir, Afikpo, Ebonyi State, Nigeria. *Animal Research International* **11**, 1958–1963.
- Ngwu, G.I., Ohaegbula, A.B.O. & Okafor, F.C. (2004) Prevalence of *Fasciola gigantica*, *Cysticercus bovis* and some other disease conditions of cattle slaughtered in Nsukka urban abattoir. *Animal Research International* **1**, 7–11.
- Njoku-Tony, R.F. (2011) Effects of some physico-chemical parameters on abundance of intermediate snails of animal trematodes in Imo State, Nigeria **3**, 15–21.
- Nnabuife, H.E., Dakul, A.D., Dogo, G.I., Egwu, O.K., Weka, P.R., Ogo, I.N., Onovoh, E.O. & Obaloto, B.O. (2013) A study on helminthiasis of cattle herds in Kachia grazing reserve of Kaduna State, Nigeria. *Veterinary World* **6**, 936–940.
- Novobilsky, A., Averpil, H.B. & Hoglund, J. (2012) The field evaluation of albendazole and triclabendazole efficacy against *Fasciola hepatica* by coproantigen ELISA in naturally infected sheep. *Veterinary Parasitology* **190**, 272–276.
- Nwigwe, J.O., Njoku, O.O., Odikamnor, O.O. & Uhuo, A.C. (2013) Comparative study of intestinal helminths and protozoa of cattle and goats in Abakaliki metropolis of Ebonyi State, Nigeria. *Advances in Applied Science Research* **4**, 223–227.
- Nwosu, C.O. & Srivastava, G.C. (1993) Liver fluke infections in livestock in Borno State, Nigeria. *Veterinary Quarterly* **15**, 182–183.
- Nzalawahe, J., Kassuku, A., Stothard, J., Coles, G. & Eisler, M. (2014) Trematode infections in cattle in Arumeru District, Tanzania are associated with irrigation. *Parasites & Vectors* **7**, 107.
- Nzalawahe, J., Kassuku, A.A., Stothard, J.R., Coles, G.C. & Eisler, M.C. (2015) Associations between trematode

- infections in cattle and freshwater snails in highland and lowland areas of Iringa Rural District, Tanzania. *Parasitology* **142**, 1430–1439.
- Odigie, B.E. & Odigie, J.O.** (2013) Fascioliasis in cattle: A survey of abattoirs in Egor, Ikpoba-Okha and Oredo Local Government Areas of Edo State, using histochemical techniques. *International Journal of Basic, Applied and Innovative Research* **2**, 1–9.
- Ogunrinade, A. & Adegoke, G.O.** (1982) Bovine fascioliasis in Nigeria – intercurrent parasitic and bacterial infections. *Tropical Animal Health and Production* **14**, 121–125.
- Ogunrinade, A. & Ogunrinade, B.I.** (1980) Economic importance of bovine fascioliasis in Nigeria. *Tropical Animal Health and Production* **12**, 155–160.
- Okaiyeto, S.O., Salami, O.S., Danbirni, S.A., Allam, L. & Onoja, I.I.** (2012) Clinical, gross and histopathological changes associated with chronic fasciolosis infection in a dairy farm. *Journal of Veterinary Advances* **2**, 444–448.
- Ollerenshaw, C.B.** (1971) Some observation on the epidemiology of fasciolosis in relation to timing of molluscicide application in the control of the disease. *Veterinary Records* **88**, 152–164.
- Omowaye, S.O., Idachaba, O.S. & Falola, O.O.** (2012) The prevalence of parasitic infections in bile of cattle slaughtered in Jos abattoir, Plateau State, Nigeria. *Global Journal of Bio-science and Biotechnology* **1**, 121–123.
- Omoleye, S.O., Qasim, A.M., Olugbon, A.S., Adu, O.A., Adam, Y.V. & Joachim, C.O.** (2012) Fasciolosis in slaughtered cattle from abattoirs in Ondo State, Nigeria. *Vom Journal of Veterinary Sciences* **9**, 47–53.
- Ondriska, F., Sobota, K., Janosek, J. & Joklova, E.** (1989) A rare case of human autochthonous dicrocoeliasis in Czechoslovakia. *Bratislavske lekarske listy* **90**, 467–469.
- Onyeabor, A.** (2014) Prevalence of bovine fasciolosis observed in three major abattoirs in Abia State Nigeria. *Journal of Veterinary Advances* **4**, 752–755.
- Opara, K.N.** (2005) Population dynamics of *Fasciola gigantica* in cattle slaughtered in Uyo, Nigeria. *Tropical Animal Health and Production* **37**, 363–368.
- Otranto, D. & Traversa, D.** (2002) A review of dicrocoeliosis of ruminants including recent advances in the diagnosis and treatment. *Veterinary Parasitology* **107**, 317–335.
- Otranto, D. & Traversa, D.** (2003) Dicrocoeliosis of ruminants: a little known fluke disease. *Trends in Parasitology* **19**, 12–15.
- Otranto, D., Rehbein, S., Weigl, S., Cantacessi, C., Parisi, A., Lia, R.P. & Olson, P.D.** (2007) Morphological and molecular differentiation between *Dicrocoelium dendriticum* (Rudolphi, 1819) and *Dicrocoelium chinensis* (Sudarikov and Ryjikov, 1951) Tang and Tang, 1978 (Platyhelminthes: Digenea). *Acta Tropica* **104**, 91–98.
- Pfukenyi, D.M., Monrad, J. & Mukaratirwa, S.** (2005) Epidemiology of trematode infections in cattle in Zimbabwe: a review. *Journal of the South African Veterinary Association* **76**, 9–17.
- Pfukenyi, D.M., Mukaratirwa, S., Willingham, A.L. & Monrad, J.** (2006a) Epidemiological studies of *Fasciola gigantica* infections in cattle in the highveld and lowveld communal grazing areas of Zimbabwe. *Onderstepoort Journal of Veterinary Research* **73**, 37–51.
- Pfukenyi, D.M., Mukaratirwa, S., Willingham, A.L. & Monrad, J.** (2006b) Epidemiological studies of *Fasciola gigantica* infections in cattle in the highveld and lowveld communal grazing areas of Zimbabwe. *Onderstepoort Journal of Veterinary Research* **73**, 179–191.
- Phiri, A.M., Phiri, I.K., Sikasunge, C.S. & Monrad, J.** (2005a) Prevalence of fasciolosis in Zambian cattle observed at selected abattoirs with emphasis on age, sex and origin. *Journal of Veterinary Medicine Series B – Infectious Diseases and Veterinary Public Health* **52**, 414–416.
- Phiri, A.M., Phiri, I.K., Siziya, S., Sikasunge, C.S., Chembensofu, M. & Monrad, J.** (2005b) Seasonal pattern of bovine fasciolosis in the Kafue and Zambezi catchment areas of Zambia. *Veterinary Parasitology* **134**, 87–92.
- Pugh, R.N., Schillhorn Van Veen, T.W. & Tayo, M.A.** (1980) Malumfashi Endemic Diseases Research Project XII. *Schistosoma bovis* and *Fasciola*. *Annals of Tropical Medicine and Parasitology* **74**, 447–453.
- Radostits, O.M., Gay, C.C., Blood, D.C. & Hinchcliff, K.W.** (2000) *Veterinary medicine: A textbook of diseases of cattle, sheep, goats, pigs and horses*. 9th edn. Philadelphia, W.B. Saunders.
- Raper, A.B.** (1951) *Schistosoma bovis* infection in man. *East African Medical Journal* **28**, 50–54.
- Roche, P.J.L.** (1948) Human dicrocoeliasis in Nigeria. *Transactions of the Royal Society of Tropical Medicine and Hygiene* **41**, 819–820.
- Rojo-Vázquez, F.A., Meana, A., Valcárcel, F. & Martínez-Valladares, M.** (2012) Update on trematode infections in sheep. *Veterinary Parasitology* **189**, 15–38.
- Rolfe, P.F. & Boray, J.C.** (1987) Chemotherapy of paramphistomosis in cattle. *Australian Veterinary Journal* **64**, 328–332.
- Rolfe, P.F., Boray, J.C., Nichols, P. & Collins, G.H.** (1991) Epidemiology of paramphistomosis in cattle. *International Journal for Parasitology* **21**, 813–819.
- Samaila, M.O., Shehu, S.M., Abubakar, N., Mohammed, U. & Jabo, B.** (2009) Human dicrocoeliasis presenting as a subcutaneous mass. *BMJ Case Reports* bcr0220091622.
- Sanabria, R., Ceballos, L., Moreno, L., Romero, J., Lanusse, C. & Alvarez, L.** (2013) Identification of a field isolate of *Fasciola hepatica* resistant to albendazole and susceptible to triclabendazole. *Veterinary Parasitology* **193**, 105–110.
- Sandoval, H., Manga-González, Y., Campo, R., García, P., Castro, J.M.A. & De La Vega, M.P.** (1999) Preliminary study on genetic variability of *Dicrocoelium dendriticum* determined by random amplified polymorphic DNA. *Parasitology International* **48**, 21–26.
- Santoro, F.** (1988) *Schistosoma bovis* in human stools in Republic of Niger. *Transactions of the Royal Society of Tropical Medicine and Hygiene* **82**, 257.
- Savioli, L., Chistulo, L. & Montresor, A.** (1999) New opportunities for the control of fasciolosis. *Bulletin World Health Organization* **77**, 300.
- Schillhorn van Veen, T.W.** (1980) Fascioliasis (*Fasciola gigantica*) in West Africa: a review. *Veterinary Bulletin* **50**, 529–533.
- Schillhorn van Veen, T.W., Shonekan, R.A.O. & Fabiyi, J.P.** (1975) A host–parasite checklist of helminth

- parasites of domestic animals in Northern Nigeria. *Bulletin of Animal Health and Production in Africa* **23**, 269–288.
- Schillhorn van Veen, T.W., Folaranmi, D.O.B., Usman, S. & Ishaya, T. (1980) Incidence of liver fluke infections (*Fasciola gigantica* and *Dicrocoelium hospes*) in ruminants in northern Nigeria. *Tropical Animal Health and Production* **12**, 97–104.
- Sewell, M.M.H. (1966) The pathogenesis of fascioliasis. *British Veterinary Association Annual Congress* **78**, 98–105.
- Sisay, A. & Nibret, E. (2013) Prevalence and risk factors of bovine and ovine fasciolosis, and evaluation of direct sedimentation sensitivity method at Bahir-Dar Municipal Abattoir, Northern Ethiopia. *Ethiopia Veterinary Journal* **17**, 1–17.
- Soulsby, E.J.L. (1982) *Helminths, arthropods and protozoa of domesticated animals*. London, Baillière Tindall.
- Spence, S.A., Fraser, G.C. & Chang, S. (1996) Responses in milk production to the control of gastrointestinal nematode and paramphistome parasites in dairy cattle. *Australian Veterinary Journal* **74**, 456–459.
- Spithill, T.W., Smooker, P.M. & Copeman, D.B. (1999) *Fasciola gigantica*: epidemiology, immunology and molecular biology. pp. 465–525 in Dalton, J.P. (Ed.) *Fasciolosis*. Wallingford, CABI Publishing.
- Stemmermann, G.N. (1953) Human infestation with *Fasciola gigantica*. *American Journal of Pathology* **29**, 731–759.
- Sugun, S.Y., Ehizibolo, D.O., Ogo, N.I., Timothy, S.Y. & Ngulukun, S.S. (2010) Prevalence of bovine fasciolosis in Bauchi State, Nigeria. *Sahel Journal of Veterinary Science* **9**, 16–20.
- Taylor, E.L. (1964) *Fasciolosis and the liver fluke*. Rome, Italy, Food and Agriculture Organization of the United Nations Agricultural Studies.
- Taylor, M.A., Coop, R.L. & Wall, R.L. (2007) *Veterinary parasitology*. Somerset, New Jersey, USA, Blackwell Publishing.
- Titi, A., Mekroud, A., Sedraoui, S., Vignoles, P. & Rondelaud, D. (2010) Prevalence and intensity of *Paramphistomum daubneyi* infections in cattle from north-eastern Algeria. *Journal of Helminthology* **84**, 177–181.
- Toledo, R., Esteban, J.G. & Fried, B. (2012) Current status of food-borne trematode infections. *European Journal of Clinical Microbiology and Infectious Diseases* **31**, 1705–1718.
- Toolan, D.P., Mitchell, G., Searle, K., Sheehan, M., Skuce, P.J. & Zadoks, R.N. (2015) Bovine and ovine rumen fluke in Ireland – prevalence, risk factors and species identity based on passive veterinary surveillance and abattoir findings. *Veterinary Parasitology* **212**, 168–174.
- Torgerson, P. & Claxton, J. (1999) Epidemiology and control. pp. 113–149 in Dalton, J.P. (Ed.) *Fasciolosis*. Wallingford, CABI Publishing.
- Ulayi, B.M., Umaru-Sule, B. & Adamu, S. (2007) Prevalence of *Dicrocoelium hospes* and *Fasciola gigantica* infections in cattle at slaughter in Zaria, Nigeria. *Journal of Animal and Veterinary Advances* **6**, 1112–1115.
- Umar, A.G., Nwosu, C.O. & Philip, H.R. (2009) Seasonal prevalence and economic importance of bovine fascioliasis in Jalingo Abattoir, Taraba State, Nigeria. *Nigerian Veterinary Journal* **30**, 44–50.
- Velusamy, R., Singh, B.P. & Raina, O.K. (2004) Detection of *Fasciola gigantica* infection in snails by polymerase chain reaction. *Veterinary Parasitology* **120**, 85–90.
- Vercruysse, J. & Gabriel, S. (2005) Immunity to schistosomiasis in animals, an update. *Parasite Immunology* **27**, 289–295.
- Walker, S.M., Makundi, A.E., Namuba, F.V., Kassuku, A.A., Keyyu, J., Hoey, E.M., Prodohl, P., Stothard, J.R. & Trudgett, A. (2008) The distribution of *Fasciola hepatica* and *Fasciola gigantica* within southern Tanzania – constraints associated with the intermediate host. *Parasitology* **135**, 495–503.
- Wanyangu, S.W., Bain, R.K., Rugutt, M.K., Nginyi, J.M. & Mugambi, J.M. (1996) Anthelmintic resistance amongst sheep and goats in Kenya. *Preventive Veterinary Medicine* **25**, 285–290.
- Webster, B.L., Rollinson, D., Stothard, J.R. & Huyse, T. (2010) Rapid diagnostic multiplex PCR (RD-PCR) to discriminate *Schistosoma haematobium* and *S. bovis*. *Journal of Helminthology* **84**, 107–114.
- WHO (1995) *Control of foodborne trematode infections*. WHO Technical Report Series. Geneva, Switzerland, World Health Organization.
- Winkelhagen, A.J.S., Mank, T., De Vries, P.J. & Soetekouw, R. (2012) Apparent triclabendazole-resistant human *Fasciola hepatica* infection, the Netherlands. *Emerging Infectious Diseases* **18**, 1028–1029.
- Wolfe, M.S. (1966) Spurious infection with *Dicrocoelium hospes* in Ghana. *American Journal of Tropical Medicine and Hygiene* **15**, 180–182.
- Wright, C.A. (1971) *Flukes and snails*. London, George Allen and Unwin.
- Yabe, J., Phiri, I.K., Phiri, A.M., Chembensofu, M., Dorny, P. & Vercruysse, J. (2008) Concurrent infections of *Fasciola*, *Schistosoma* and *Amphistomum* spp. in cattle from Kafue and Zambezi river basins of Zambia. *Journal of Helminthology* **82**, 373–376.
- Yahaya, A. & Tyav, Y.B. (2014) A survey of gastrointestinal parasitic helminths of bovine slaughtered in abattoir, Wudil Local Government Area, Kano State, Nigeria. *Greener Journal of Biological Sciences* **4**, 128–134.
- Yilma, J.M. & Mesfin, A. (2000) Dry season bovine fasciolosis in Northwestern part of Ethiopia. *Revue de Medecine Veterinaire* **6**, 493–500.
- Zintl, A., Garcia-Campos, A., Trudgett, A., Chrysafidis, A.L., Talavera-Arce, S., Fu, Y., Egan, S., Lawlor, A., Negro, C., Brennan, G., Hanna, R.E., De Waal, T. & Mulcahy, G. (2014) Bovine paramphistomes in Ireland. *Veterinary Parasitology* **204**, 199–208.