

Article

Inauguration of Kifilideen theorem of matrix transformation of negative power of $-n$ of trinomial expression in which three variables x , y and z are found in parts of the trinomial expression with some other applications

Kifilideen L. Osanyinpeju

Agricultural and Bio-Resources Engineering Department, College of Engineering, Federal University of Agriculture Abeokuta, Ogun State, Nigeria

E-mail: amkifilideenosanyinpeju@gmail.com, prof_4us@yahoo.com

Received 27 November 2021



Abstract

Kifilideen trinomial theorem of negative power of $-n$ is theorem which is used to generate the series and terms of a trinomial expression of negative power of $-n$ in an orderly and periodicity manner that is based on standardized and matrix methods. Negative power of Newton binomial theorem had been used to produce series of partial fractions of a compound fraction. The establishment of the negative power of $-n$ of trinomial theorem would extend the number of compound fraction in which series (expansion) can be produced. This study applied Kifilideen expansion of negative power of $-n$ of Kifilideen trinomial theorem for the transformation of compound fraction into series of partial fractions with other developments. A theorem of matrix transformation of negative power of $-n$ of trinomial expression in which three variables x , y and z are found in parts of the trinomial expression was inauguration. The development would ease the process of evaluating such trinomial expression of negative power of $-n$. This standardized and matrix method used in arranging the terms of the Kifilideen expansion of negative power of $-n$ of trinomial expression yield an interesting results in which it is utilized in transforming compound fraction into series of partial fractions in a unique way.

Keywords compound fraction; series; partial fraction; combination; kif matrix; Kifilideen standardized method.

Selforganizology

ISSN 2410-0080

URL: <http://www.iaees.org/publications/journals/selforganizology/online-version.asp>

RSS: <http://www.iaees.org/publications/journals/selforganizology/rss.xml>

E-mail: selforganizology@iaees.org

Editor-in-Chief: WenJun Zhang

Publisher: International Academy of Ecology and Environmental Sciences

1 Introduction

Sir Isaac Newton (1642–1727) was the first mathematician and physicist to inaugurate negative and fraction power of binomial theorem (Dennis and Addington, 2009; Goss, 2011; Youngmee and Sangwook, 2014;

Schwartz, 2015; Cooley, 2019; Osanyinpeju, 2019a). He established formula for binomial theorem that could work for negative, fraction powers of binomial expression (Francia, 2000; Haggstrom, 2012; Aljohani, 2016; Anekwe, 2018; Tavora, 2020). The binomial theorem is a general expression for any power of the sum or difference of any two things, terms or quantities (Godman et al., 1984, Talber et al., 1995; Bird, 2003; Stroud and Booth, 2007, Tuttuh et al., 2014; Bunday and Mulholland, 2014). Binomial is widely used in the field of Physics, Biology and Engineering to expand power of binomial expression (Costa, 2017; Gavrikov, 2018). Binomial series (expansions) are utilized in area of mathematical analysis link to data modeling, probability theory, algebra and approximation techniques (Yang, 2017).

Kifilideen trinomial theorem of negative power of $-n$ is theorem which is used to generate the series and terms of a trinomial expression of negative power of $-n$ in an orderly and periodicity manner that is based on standardized and matrix method (Osanyinpeju, 2021a, 2021b). Negative power of Newton binomial theorem had been used to produce series of partial fractions of a compound fraction (Horn, 2020). The establishment of the negative power of $-n$ of trinomial theorem would extend the number of compound fraction in which series can be produced. The negative power of $-n$ of Kifilideen trinomial theorem can be used to generate the series of compound fraction number (examples are partial fraction series of $\frac{13}{9}, \frac{1}{1.75}, \frac{729}{811}$).

The terms of negative power of $-n$ of trinomial theorem produce an infinite series unlike the terms of power of n of trinomial theorem which generate finite series (Osanyinpeju, 2020a). The terms of negative power of $-n$ of trinomial theorem when arranged in the form of matrix, the number of elements in each column has finite value which is also the same for positive power of n of trinomial theorem. Although, the column of any negative power of $-n$ of trinomial theorem is infinite while that of positive power of n of trinomial theorem is finite.

The power combination of each term of the negative power of $-n$ of Kifilideen trinomial theorem is arranged in groups (columns) and periods (rows) in the kif matrix where each power combination takes a designated position. No two power combinations have the same position in the kif matrix for a particular negative power of $-n$ of Kifilideen trinomial theorem. This is also applied to positive power of trinomial theorem (Osanyinpeju, 2020b). The power combination of each term; down the group (column) and across the period (row) decreases and increases respectively at regular pattern for both positive and negative powers of Kifilideen trinomial theorem in the kif matrix format. Also, to transit from one group to another the power combination progress with unique addition of regular figure which occur for both positive and negative power of trinomial theorem.

This standardized and matrix method used in arranging the terms of the Kifilideen expansion of negative power of $-n$ of trinomial expression yield an interesting results in which it is utilized in transforming compound fraction into series of partial fractions in a unique way. Kifilideen Matrix approach had been extended in evaluating and computation of power of base of eleven, other bi-digits and tri-digit numbers (Osanyinpeju, 2019b, 2020c, 2020d). This study applied Kifilideen expansion of negative power of $-n$ of Kifilideen trinomial theorem for the transformation of compound fraction into series of partial fraction with other development.

2 Materials and Methods

The Kifilideen coefficient table of negative power of Kifilideen trinomial theorem was used in generating the power coefficient of each term of series of the trinomial expansion.

2.1 Application of Kifilideen series (expansion) of negative power of – n of trinomial theorem using standardized method

[a] Generate the series of $[1 + x + x^3]^{-1}$ using Kifilideen standardized method. Otherwise determine the series of $\frac{343}{393}$ (Hint: take $x = \frac{1}{7}$).

Solution

[a] Using Kifilideen coefficient table of negative power of – 1 of trinomial theorem, the coefficient in ascending order 1, - 1, - 1, 1, 2, 1, - 1, - 3, - 3, - 1, 1, 4, 6, 4, 1, -1, - 5, - 10, - 10, - 5, - 1, 1, 6, 15, 20, 15, 6, 1, ...

$$\begin{aligned}
 [1 + x + x^3]^{-1} &= 1 - x^1[x^3]^0 - x^0[x^3]^1 + x^2[x^3]^0 + 2x^1[x^3]^1 + x^0[x^3]^2 - x^3[x^3]^0 - 3x^2[x^3]^1 - 3x^1[x^3]^2 - x^0[x^3]^3 + \\
 &x^4[x^3]^0 + 4x^3[x^3]^1 + 6x^2[x^3]^2 + 4x^1[x^3]^3 + x^0[x^3]^4 - x^5[x^3]^0 - 5x^4[x^3]^1 - 10x^3[x^3]^2 - 10x^2[x^3]^3 - 5x^1[x^3]^4 \\
 &- x^0[x^3]^5 + x^6[x^3]^0 + 6x^5[x^3]^1 + 15x^4[x^3]^2 + 20x^3[x^3]^3 + 15x^2[x^3]^4 + 6x^1[x^3]^5 + x^0[x^3]^6 \\
 &- x^7[x^3]^0 - 7x^6[x^3]^1 - \\
 &21x^5[x^3]^2 - 35x^4[x^3]^3 - 35x^3[x^3]^4 - 21x^2[x^3]^5 - 7x^1[x^3]^6 - x^0[x^3]^7 + x^8[x^3]^0 + 8x^7[x^3]^1 + 28x^6[x^3]^2 + 56x^5[x^3]^3 \\
 &+ 70x^4[x^3]^4 + 56x^3[x^3]^5 + 28x^2[x^3]^6 + 8x^1[x^3]^7 + x^0[x^3]^8 - x^9[x^3]^0 - 9x^8[x^3]^1 - 36x^7[x^3]^2 - 84x^6[x^3]^3 - \\
 &126x^5[x^3]^4 - 126x^4[x^3]^5 - 84x^3[x^3]^6 - 36x^2[x^3]^7 - 9x^1[x^3]^8 - x^{10}[x^3]^0 + 10x^9[x^3]^1 + 45x^8[x^3]^2 + 120x^7[x^3]^3 \\
 &+ 210x^6[x^3]^4 + 252x^5[x^3]^5 + 210x^4[x^3]^6 + 120x^3[x^3]^7 + 45x^2[x^3]^8 + 10x^1[x^3]^9 + x^0[x^3]^{10} - x^{11}[x^3]^0 - \dots \quad [1]
 \end{aligned}$$

$$\begin{aligned}
 [1 + x + x^3]^{-1} &= 1 - x - x^3 + x^2 + 2x^4 + x^6 - x^3 - 3x^5 - 3x^7 - x^9 + x^4 + 4x^6 + 6x^8 + 4x^{10} + x^{12} - x^5 - 5x^7 - 10x^9 \\
 &- 10x^{11} - 5x^{13} - x^{15} + x^6 + 6x^8 + 15x^{10} + 20x^{12} + 15x^{14} + 6x^{16} + x^{18} - x^7 - 7x^9 - 21x^{11} - 35x^{13} - 35x^{15} - 21x^{17} \\
 &- 7x^{19} - x^{21} + x^8 + 8x^{10} + 28x^{12} + 56x^{14} + 70x^{16} + 56x^{18} + 28x^{20} + 8x^{22} + x^{24} - x^7 - 9x^{11} - 36x^{13} - 84x^{15} - \\
 &126x^{17} - 126x^{19} - 84x^{21} - 36x^{23} - 9x^{25} - x^{27} + x^{10} + 10x^{12} + 45x^{14} + 120x^{16} + 210x^{18} + 252x^{20} + 210x^{22} + \\
 &120x^{24} + 45x^{26} + 10x^{28} + x^{30} - x^{11} - \dots \quad [2]
 \end{aligned}$$

$$[1 + x + x^3]^{-1} = 1 - x + x^2 - 2x^3 + 3x^4 - 4x^5 + 6x^6 - 9x^7 + 13x^8 - 19x^9 + 28x^{10} - 41x^{11} + 59x^{12} - \dots$$

$$[ii] \frac{343}{393} = [1 + \frac{1}{7} + (\frac{1}{7})^3]^{-1} = 1 - \frac{1}{7} + \frac{1}{49} - \frac{2}{343} + \frac{3}{2401} - \frac{4}{16807} + \frac{6}{117649} - \frac{9}{823543} + \frac{13}{5764801} - \frac{19}{50353607} + \frac{28}{282475249} -$$

$$\frac{41}{1977326743} + \frac{59}{13841287200} - \dots \quad [3]$$

The evaluation of the above series gives 0.872774 to 6 decimal places. Also, using calculator $\frac{343}{393}$ gives 0.872774 to 6 decimal places. This indicates that the negative power of -1 of Kifilideen trinomial theorem and coefficient of negative power of -1 from the Kifilideen coefficient table are valid.

[b] Produce the Kifilideen expansion of $[3 - x + x^4]^{-2}$ using Kifilideen standardized method. Use the expansion to generate the series of $\frac{6561}{47089}$ (Hint: take $x = \frac{1}{3}$)

Solution

[b] Using Kifilideen coefficient table of negative power of -2 of trinomial theorem, the coefficient in ascending order 1, -2, -2, 3, 6, 3, -4, -12, -12, -4, 5, 20, 30, 20, 5, -6, -30, -60, -60, -30, -6, 7, 42, 105, 140, 105, 42, 7, -8, -56, -168, -280, -280, -168, -56, -8, ...

$$\begin{aligned}
 [3 - x + x^4]^{-2} &= [3]^{-2}[-x]^0[x^4]^0 - 2[3]^{-3}[-x]^1[x^4]^0 - 2[3]^{-3}[-x]^0[x^4]^1 + 3[3]^{-4}[-x]^2[x^4]^0 + 6[3]^{-4}[-x]^1[x^4]^1 + \\
 &3[3]^{-4}[-x]^0[x^4]^2 - 4[3]^{-5}[-x]^3[x^4]^0 - 12[3]^{-5}[-x]^2[x^4]^1 - 12[3]^{-5}[-x]^1[x^4]^2 - 4[3]^{-5}[-x]^0[x^4]^3 + 5[3]^{-6}[-x]^4[x^4]^0 \\
 &+ 20[3]^{-6}[-x]^3[x^4]^1 + 30[3]^{-6}[-x]^2[x^4]^2 + 20[3]^{-6}[-x]^1[x^4]^3 + 5[3]^{-6}[-x]^0[x^4]^4 - 6[3]^{-7}[-x]^5[x^4]^0 \\
 &- 30[3]^{-7}[-x]^4[x^4]^1 - 60[3]^{-7}[-x]^3[x^4]^2 - 60[3]^{-7}[-x]^2[x^4]^3 - 30[3]^{-7}[-x]^1[x^4]^4 - 6[3]^{-7}[-x]^0[x^4]^5 + \\
 &7[3]^{-8}[-x]^6[x^4]^0 + 42[3]^{-8}[-x]^5[x^4]^1 + 105[3]^{-8}[-x]^4[x^4]^2 + 140[3]^{-8}[-x]^3[x^4]^3 + 105[3]^{-8}[-x]^2[x^4]^4 \\
 &+ 42[3]^{-8}[-x]^1[x^4]^5 + 7[3]^{-8}[-x]^0[x^4]^6 - \dots
 \end{aligned} \tag{4}$$

$$\begin{aligned}
 [3 - x + x^4]^{-2} &= \frac{1}{9} + \frac{2x}{27} - \frac{2x^4}{27} + \frac{3x^2}{81} - \frac{6x^5}{81} + \frac{3x^8}{81} + \frac{4x^3}{243} - \frac{12x^6}{243} + \frac{12x^9}{243} - \frac{4x^{12}}{243} + \frac{5x^4}{729} - \frac{20x^7}{729} + \frac{30x^{10}}{729} - \frac{20x^{13}}{729} + \frac{5x^{16}}{729} \\
 &+ \frac{6x^5}{2187} - \frac{30x^8}{2187} + \\
 &\frac{60x^{15}}{2187} - \frac{60x^{14}}{2187} + \frac{30x^{17}}{2187} - \frac{6x^{20}}{2187} + \dots
 \end{aligned} \tag{5}$$

$$\begin{aligned}
 [3 - x + x^4]^{-2} &= \frac{1}{9} + \frac{2x}{27} + \frac{3x^2}{81} + \frac{4x^3}{243} - \frac{2x^4}{27} + \frac{5x^4}{729} - \frac{6x^5}{81} + \frac{6x^5}{2187} - \frac{12x^6}{243} - \frac{20x^7}{729} + \frac{3x^8}{81} - \frac{30x^8}{2187} + \frac{12x^9}{243} + \frac{30x^{10}}{729} - \frac{4x^{12}}{243} \\
 &- \frac{20x^{13}}{729} - \\
 &\frac{60x^{14}}{2187} + \frac{60x^{15}}{2187} + \frac{5x^{16}}{729} + \frac{30x^{17}}{2187} - \frac{6x^{20}}{2187} + \dots
 \end{aligned} \tag{6}$$

$$[ii] \frac{6561}{47089} = \left[3 - \frac{1}{3} + \left(\frac{1}{3} \right)^4 \right]^{-2} = \frac{1}{9} + \frac{2}{81} + \frac{3}{729} + \frac{4}{6561} - \frac{2}{2187} + \frac{5}{59049} - \frac{6}{19683} + \frac{6}{531441} - \frac{12}{177147} - \frac{20}{1594323} + \frac{3}{531441} - \frac{30}{14348907} + \frac{12}{4782969} + \frac{30}{43046721} - \frac{4}{129140163} \dots \quad [7]$$

The evaluation of the above series gives 0.13933 to 5 decimal places. Also, using calculator $\frac{6561}{47089}$ gives 0.13933 to 5 decimal places. This indicates that the negative power of - 2 of Kifilideen trinomial theorem and coefficient of negative power of - 2 from the Kifilideen coefficient table are valid.

2.2 Generating series of partial fractions of compound fraction which denominator is decimal number

[d] Produce the series of $[1 + x + x^2]^{-1}$ using Kifilideen standardized method. Otherwise determine the series of

$$\frac{1}{1.75} \text{ (Hint: } x = \frac{1}{2} \text{)}.$$

Solution

Using Kifilideen coefficient table of negative power of - 1 of trinomial theorem, the coefficient in ascending order 1, - 1, - 1, 1, 2, 1, - 1, - 3, - 3, - 1, 1, 4, 6, 4, 1, -1, - 5, - 10, - 10, - 5, - 1, 1, 6, 15, 20, 15, 6, 1, ...

$$\begin{aligned} [1 + x + x^2]^{-1} &= 1 - x^1[x^2]^0 - x^0[x^2]^1 + x^2[x^2]^0 + 2x^1[x^2]^1 + x^0[x^2]^2 - x^3[x^2]^0 - 3x^2[x^2]^1 - 3x^1[x^2]^2 - x^0[x^2]^3 + \\ &x^4[x^2]^0 + 4x^3[x^2]^1 + 6x^2[x^2]^2 + 4x^1[x^2]^3 + x^0[x^2]^4 - x^5[x^2]^0 - 5x^4[x^2]^1 - 10x^3[x^2]^2 - 10x^2[x^2]^3 - 5x^1[x^2]^4 - \\ &x^0[x^2]^5 + x^6[x^2]^0 + 6x^5[x^2]^1 + 15x^4[x^2]^2 + 20x^3[x^2]^3 + 15x^2[x^2]^4 + 6x^1[x^2]^5 + x^0[x^2]^6 - x^7[x^2]^0 - 7x^6[x^2]^1 - \\ &21x^5[x^2]^2 - 35x^4[x^2]^3 - 35x^3[x^2]^4 - 21x^2[x^2]^5 - 7x^1[x^2]^6 - x^0[x^2]^7 + x^8[x^2]^0 + 8x^7[x^2]^1 + 28x^6[x^2]^2 + 56x^5[x^2]^3 \\ &+ 70x^4[x^2]^4 + 56x^3[x^2]^5 + 28x^2[x^2]^6 + 8x^1[x^2]^7 + x^0[x^2]^8 - x^9[x^2]^0 - 9x^8[x^2]^1 - 36x^7[x^2]^2 - 84x^6[x^2]^3 - \\ &126x^5[x^2]^4 - 126x^4[x^2]^5 - 84x^3[x^2]^6 - 36x^2[x^2]^7 - 9x^1[x^2]^8 - x^10[x^2]^0 + 10x^9[x^2]^1 + 45x^8[x^2]^2 + 120x^7[x^2]^3 + \\ &210x^6[x^2]^4 + 252x^5[x^2]^5 + 210x^4[x^2]^6 + 120x^3[x^2]^7 + 45x^2[x^2]^8 + 10x^1[x^2]^9 + x^0[x^2]^10 - x^{11}[x^2]^0 - \dots \quad [8] \end{aligned}$$

$$\begin{aligned} [1 + x + x^3]^{-1} &= 1 - x - x^2 + x^2 + 2x^3 + x^4 - x^3 - 3x^4 - 3x^5 - x^6 + x^4 + 4x^5 + 6x^6 + 4x^7 + x^8 - x^5 - 5x^6 - 10x^7 \\ &- 10x^8 - 5x^9 - x^{10} + x^6 + 6x^7 + 15x^8 + 20x^9 + 15x^{10} + 6x^{11} + x^{12} - x^7 - 7x^8 - 21x^9 - 35x^{10} - 35x^{11} - 21x^{12} - \\ &7x^{13} - x^{14} + x^8 + 8x^9 + 28x^{10} + 56x^{11} + 70x^{12} + 56x^{13} + 28x^{14} + 8x^{15} + x^{16} - x^9 - 9x^{10} - 36x^{11} - 84x^{12} - 126x^{13} \\ &- 126x^{14} - 84x^{15} - 36x^{16} - 9x^{17} - x^{18} + x^{10} + 10x^{11} + 45x^{12} + 120x^{13} + 210x^{14} + 252x^{15} + 210x^{16} + 120x^{17} + \end{aligned}$$

$$\begin{aligned}
&45x^{18} + 10x^{19} + x^{20} - x^{11} - 11x^{12} - 55x^{13} - 165x^{14} - 330x^{15} - 462x^{16} - 462x^{17} - 330x^{18} - 165x^{19} - 55x^{20} - \\
&11x^{21} - x^{22} + x^{12} + 12x^{13} + 66x^{14} + 220x^{15} + 495x^{16} + 792x^{17} + 924x^{18} + 792x^{19} + 495x^{20} + 220x^{21} + 66x^{22} + \\
&12x^{23} + x^{24} - x^{13} + \dots
\end{aligned}$$

[9]

$$[1 + x + x^3]^{-1} = 1 - x + x^3 - x^4 + x^6 - x^7 + x^9 - x^{10} + x^{12} - x^{13} \dots$$

[10]

$$[ii] \frac{1}{1.75} = [1 + 0.5 + (0.5)^2]^{-1} = 1 - \frac{1}{2} + \frac{1}{8} - \frac{1}{16} + \frac{1}{64} - \frac{1}{128} + \frac{1}{512} - \frac{1}{1024} + \frac{1}{4096} - \frac{1}{8192} + \dots$$

[11]

The evaluation of the above series gives 0.5714 to 4 decimal places. Also, using calculator $\frac{1}{1.75}$ gives 0.5714 to 4 decimal places. This indicates that the negative power of -1 of Kifilideen trinomial theorem and coefficient of negative power of -1 from the Kifilideen coefficient table are valid.

2.3 General application of negative power of – n of Kifilideen trinomial theorem using standardized method

[e] If a t^{th} term in the Kifilideen expansion of $[1 - 2x + ay^2]^{-3}$ yield $-7680xy^6$. Determine the following:

[i] the power combination of the term

[ii] the t^{th} term

[iii] the value of a

Solution

[i] For any given term in the Kifilideen expansion of $[1 - 2x + ay^2]^{-3}$ is given as

$${}_{kif}^{-3}C [1]^k [-2x]^i [ay^2]^f$$

[12]

$${}_{kif}^{-3}C [1]^k [-2]^i [x]^i [a]^f [y^2]^f$$

[13]

Comparing [13] with $-7680xy^6$, so

$$i = 1$$

[14]

$$2f = 6$$

[15]

$$\text{Therefore, } f = 3$$

[16]

Also,

$$k + i + f = n$$

[17]

where, k, i , and f are the component parts of the power combination

n – the negative power of – n of the trinomial expression

$$k + i + f = n \quad [18]$$

$$k + 1 + 3 = -3 \quad [19]$$

$$k = -7 \quad [20]$$

$$C_p = kif = -713 \quad [21]$$

[ii]

$$t = \frac{[n - k]^2 + [n - k] + 2f + 2}{2} \quad [22]$$

$$k = -7, i = 1, f = 3 \text{ and } n = -3 \quad [23]$$

$$t = \frac{[-3 - -7]^2 + [-3 - -7] + 2 \times 3 + 2}{2} \quad [24]$$

$$t = 14^{\text{th}} \text{ term} \quad [25]$$

$$[iii] \quad {}_{kif}^{-3}C [1]^k [-2]^i [x]^i [a]^f [y^2]^f = -7680xy^6 \quad [26]$$

$${}_{-713}^{-3}C [1]^{-7} [-2]^1 [x]^1 [a]^3 [y^2]^3 = -7680xy^6 \quad [27]$$

From the Kifilideen coefficient table, the coefficient of the 14th term on the $n = -3$ column is + 60 or

$$\frac{-3!}{-7!1!3!} [1]^{-7} [-2]^1 [x]^1 [a]^3 [y^2]^3 = -7680xy^6 \quad [28]$$

$$\frac{-3 \times -4 \times -5 \times -6 \times -7!}{-7!1!3!} [1]^{-7} [-2]^1 [x]^1 [a]^3 [y^2]^3 = -7680xy^6 \quad [29]$$

$$\frac{-3 \times -4 \times -5 \times -6}{1!3!} [1]^{-7} [-2]^1 [x]^1 [a]^3 [y^2]^3 = -7680xy^6 \quad [30]$$

$$+60 \times [1]^{-7} [-2]^1 [x]^1 [a]^3 [y^2]^3 = -7680xy^6 \quad [31]$$

$$+60 \times -2 \times a^3 \times xy^6 = -7680xy^6 \quad [32]$$

$$a^3 = \frac{-7680}{-120} \quad [33]$$

$$a^3 = 64 \quad [34]$$

$$a = 4 \quad [35]$$

[f] If a term of the Kifilideen expansion of $\left[1 - \frac{bx^2}{y^4} + 2yz^3\right]^{-4}$ is $7560 x^4 y^{-6} z^6$. Find

[i] the power combination of the term

[ii] the t^{th} term

[iii] the value of b

Solution

$${}_{kif}^{-4}C [1]^k \left[-\frac{bx^2}{y^4}\right]^i [2yz^3]^f \quad [36]$$

$${}_{kif}^{-4}C [1]^k [-b]^i [x^2]^i [y^{-4}]^i [2]^f [y]^f [z^3]^f \quad [37]$$

$${}_{kif}^{-4}C [1]^k [-b]^i [2]^f [x^2]^i [y]^{-4i+f} [z^3]^f \quad [38]$$

Comparing [38] with $7560 x^4 y^{-6} z^6$, so

$$2i = 4 \quad [39]$$

$$i = 2 \quad [40]$$

$$-4i + f = -6 \quad [41]$$

$$-4 \times 2 + f = -6 \quad [42]$$

$$f = 2 \quad [43]$$

Therefore,

Also,

$$k + i + f = n \quad [44]$$

$$k + 2 + 2 = -4 \quad [45]$$

$$k = -8 \quad [46]$$

$$C_p = kif = -822 \quad [47]$$

$$[ii] \ k = -8, \ i = 2 \text{ and } f = 2 \quad [48]$$

$$\text{Note,} \quad x = n - k \quad [49]$$

$$t = y + f \text{ and } n = k + i + f \quad [50]$$

So,

$$x = -4 - -8 = 4 \quad [51]$$

$$y = \frac{x^2 + x + 2}{2} \quad [52]$$

$$y = \frac{4^2 + 4 + 2}{2} \quad [53]$$

$$y = 11 \quad [54]$$

$$t = y + f = 11 + 2 = 13^{\text{th}} \text{ term} \quad [55]$$

[iii] Comparing [38] with $7560 x^4 y^{-6} z^6$, then

$${}_{kif}^{-4}C [1]^k [-b]^i [2]^f [x^2]^i [y]^{-4i+f} [z^3]^f = 7560 x^4 y^{-6} z^6 \quad [56]$$

$${}_{-822}^{-4}C [1]^{-8} [-b]^2 [2]^2 [x^2]^2 [y]^{-4 \times 2 + 2} [z^3]^2 = 7560 x^4 y^{-6} z^6 \quad [57]$$

$${}_{-822}^{-4}C [1]^{-8} [-b]^2 [2]^2 x^4 y^{-6} z^6 = 7560 x^4 y^{-6} z^6 \quad [58]$$

From the Kifilideen coefficient table, the coefficient of the 13^{th} term on the $n = -4$ column is $+210$

$$+210 [1]^{-8} [-b]^2 [2]^2 x^4 y^{-6} z^6 = 7560 x^4 y^{-6} z^6 \quad [59]$$

$$840 b^2 = 7560 \quad [60]$$

$$b^2 = \frac{7560}{840} \quad [61]$$

$$b^2 = 9 \quad [62]$$

$$b = 3 \quad [63]$$

[g] If a term in the Kifilideen expansion of $\left[\frac{cx^2}{z^5} + 5x^3y - 3yz^2\right]^{-5}$ is $\frac{850500}{1048576} x^{-14} y^5 z^{56}$. Determine the value of

[i] the power combination

[ii] show that the negative power of $-n$ of the trinomial expression is -5

[iii] the t^{th} term

[iv] the value of c .

Solution

[i]

$${}_{kif}^{-5}C \left[\frac{cx^2}{z^5}\right]^k [5x^3y]^i [-3yz^2]^f \quad [64]$$

$${}_{kif}^{-4}C [c]^k [x^2]^k [z^{-5}]^k [5]^i [x^3]^i [y]^i [-3]^f [y]^f [z^2]^f \quad [65]$$

$${}_{kif}^{-5}C [c]^k [5]^i [-3]^f [x]^{2k+3i} [y]^{i+f} [z]^{-5k+2f} \quad [66]$$

Comparing [67] with $\frac{850500}{1048576}x^{-4}y^5z^{56}$, so

$$2k + 3i = -14 \quad [67]$$

$$i + f = 5 \quad [68]$$

$$-5k + 2f = 56 \quad [69]$$

Also,

$$k + i + f = n \quad [70]$$

$$k + i + f = -5 \quad [71]$$

Put [68] in [67]

$$2k + 3[5 - f] = -14 \quad [72]$$

From [72] and [69], then

$$2k - 3f = -29 \times 5 \quad [73]$$

$$-5k + 2f = 56 \times 2 \quad [74]$$

$$10k - 15f = -145$$

$$-10k + 4f = 112$$

$$-11f = -33 \quad [77]$$

$$f = 3 \quad [78]$$

From [68] put [78] in [79], $i + f = 5$ [79]

$$i + 3 = 5 \quad [80]$$

$$i = 2 \quad [81]$$

From [67] put [81] in [82] $2k + 3i = -14 \quad [82]$

$$2k + 3[2] = -14 \quad [83]$$

$$k = -10 \quad [84]$$

$$\text{Power combination} = kif = -10, 2, 3 \quad [85]$$

[ii] From [70], $n =$ negative power of $-n$ of the trinomial expression $= k + i + f \quad [86]$

Put [78], [81] and [84] in [86], so $n = n = -10 + 2 + 3 \quad [87]$

$$n = -5 \quad [88]$$

Proved

[iii] $t = \frac{[n-k]^2 + [n-k] + 2f + 2}{2} \quad [89]$

$$k = -10, i = 2, f = 3 \text{ and } n = -5 \quad [90]$$

$$t = \frac{[-5--10]^2 + [-5--10] + 2 \times 3 + 2}{2} \quad [91]$$

$$t = 19^{\text{th}} \text{ term} \quad [92]$$

[iv] Comparing [67] with $\frac{850500}{1048576}x^{-4}y^5z^{56}$, so

$${}_{kif}^{-5}C [c]^k [5]^i [-3]^f [x]^{2k+3i} [y]^{i+f} [z]^{-5k+2f} = \frac{850500}{1048576} x^{-4} y^5 z^{56} \quad [93]$$

$${}_{-10,2,3}^{-5}C [c]^{-10} [5]^2 [-3]^3 [x]^{-4} [y]^5 [z]^{56} = \frac{850500}{1048576} x^{-4} y^5 z^{56} \quad [94]$$

From the Kifilideen coefficient table, the coefficient of the 19th term on the $n = -5$ column is -1260

$$-1260 [c]^{-10} [5]^2 [-3]^3 = \frac{850500}{1048576} \quad [95]$$

$$c^{-10} = \frac{850500}{1048576 \times 27 \times 25 \times 1260} \quad [96]$$

$$c^{-10} = [1048576]^{-1} \quad [97]$$

$$c^{-10} = [4]^{-10} \quad [98]$$

$$c = 4 \quad [99]$$

2.4 Theorem of matrix transformation of negative power of $-n$ of trinomial expression

If three variables x , y and z are found in each part of trinomial expression of negative power of $-n$ such as

$$[sx^d y^e z^g + ux^h y^j z^l + vx^m y^r z^p]^{-n} \quad [100]$$

and the power combination of any term in the Kifilideen expansion of that kind of negative power of $-n$ of the trinomial expression is set as kif while the value of this term is designated as $wx^ay^bz^c$.

Then, the Kifilideen matrix transformation of such negative power of $-n$ of trinomial expression is of the form

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix}: \quad \begin{bmatrix} d & h & m \\ e & j & r \\ g & l & p \end{bmatrix} \begin{bmatrix} k \\ i \\ f \end{bmatrix} = \begin{bmatrix} a \\ b \\ c \end{bmatrix} \quad [101]$$

Thus $k + i + f = n$ and where s, u, v and w are constants

More so,

$${}^{-n}_{kif}Cs^k u^i v^f = w \quad [102]$$

Illustration

Given that a term in the Kifilideen expansion of $\left[\frac{dx^5}{z^3} + \frac{z^7}{x^2y} + \frac{8y}{z} \right]^{-n}$ is $\frac{-z^{16}}{32x^7y}$ where d is a constant value. Using

Kifilideen matrix transformation method of negative power of $-n$ of trinomial expression. Find

[i] the power combination

[ii] the power of the trinomial expression

[iii] the t^{th} term

[iv] the value of d .

Solution

$$[i] \text{ Trinomial expression: } [dx^5y^0z^{-3} + x^{-2}y^{-1}z^7 + 8x^0y^1z^{-1}]^{-n} \quad [103]$$

$$\text{Power combination to be obtained: } k \quad i \quad f \quad [104]$$

$$t^{\text{th}} \text{ term of the power combination: } \frac{-1}{32}x^{-17}y^{-1}z^{16} \quad [105]$$

Using the Kifilideen matrix transformation method, so

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix}: \quad \begin{bmatrix} 5 & -2 & 0 \\ 0 & -1 & 1 \\ -3 & 7 & -1 \end{bmatrix} \begin{bmatrix} k \\ i \\ f \end{bmatrix} = \begin{bmatrix} -17 \\ -1 \\ 16 \end{bmatrix} \quad [106]$$

$$\text{Also, } k + i + f = n \quad [107]$$

Using Crammer's rule, so

$$k = \frac{\Delta k}{\Delta} = \frac{\begin{vmatrix} -17 & -2 & 0 \\ -1 & -1 & 1 \\ 16 & 7 & -1 \end{vmatrix}}{\begin{vmatrix} 5 & -2 & 0 \\ 0 & -1 & 1 \\ -3 & 7 & -1 \end{vmatrix}} \quad [108]$$

$$k = -3 \quad [109]$$

$$i = \frac{\Delta i}{\Delta} = \frac{\begin{vmatrix} 5 & -17 & 0 \\ 0 & -1 & 1 \\ -3 & 16 & -1 \end{vmatrix}}{\begin{vmatrix} 5 & -2 & 0 \\ 0 & -1 & 1 \\ -3 & 7 & -1 \end{vmatrix}} \quad [110]$$

$$i = 1 \quad [111]$$

$$f = \frac{\Delta f}{\Delta} = \frac{\begin{vmatrix} 5 & -2 & -17 \\ 0 & -1 & -1 \\ -3 & 7 & 16 \end{vmatrix}}{\begin{vmatrix} 5 & -2 & 0 \\ 0 & -1 & 1 \\ -3 & 7 & -1 \end{vmatrix}} \quad [112]$$

$$f = 0 \quad [113]$$

$$\text{So, the power combination} = kif = -310 \quad [114]$$

$$[\text{ii}] \text{ the negative power of } -n \text{ of the trinomial expression} = n = k + i + f = -3 + 1 + 0 \quad [115]$$

$$n = -2 \quad [116]$$

$$[\text{iii}] \quad t = \frac{[n-k]^2 + [n-k] + 2f + 2}{2} \quad [117]$$

$$t = \frac{[-2--3]^2 + [-2--3] + 2 \times 0 + 2}{2} \quad [118]$$

$$t = 2^{\text{nd}} \text{ term} \quad [119]$$

$$[\text{iv}] \text{ from the question, } s = d, u = 1, v = 8 \text{ and } w = \frac{-1}{32} \quad [120]$$

Using Kifilideen matrix transformation method,

$${}_{kif}^{-n} C s^k u^i v^f = w \quad [121]$$

$${}_{-310}^{-2} C [d]^{-3} [1]^1 [8]^0 = \frac{-1}{32} \quad [122]$$

From the Kifilideen coefficient table, the coefficient of the 2^{nd} term on the $n = -2$ column is -2

$$-2 \times [d]^{-3} = \frac{-1}{32} \quad [123]$$

$$[d]^{-3} = [64]^{-1} \quad [124]$$

$$[d]^{-3} = [4]^{-3} \quad [125]$$

$$d = 4 \quad [126]$$

3 Conclusion

This study applied Kifilideen expansion of negative power of $-n$ of Kifilideen trinomial theorem for the transformation of compound fraction into series of partial fraction with other development. A theorem of matrix transformation of negative power of $-n$ of trinomial expression in which three variables x , y and z are found in part of the trinomial expression was developed. The development would ease the process of evaluating such trinomial expression of negative power of $-n$. The standardized and matrix method used in arranging the terms of the Kifilideen expansion of negative power of $-n$ of trinomial expression yield an interesting results in which it is utilized in transforming compound fraction into series of partial fractions in a unique way.

References

- Aljohani S. 2016. History of binomial theory. *International Journal of Scientific and Engineering Research*, 7(4): 161-162
- Anekwe UO. 2018. Awekwe's corrections on the negative binomial expansion. *Scientific Review*, 4(4): 26-33
- Bird J. 2003. *Engineering Mathematics*. Newnes, Oxford, An imprint of Elsevier Science Linacre House, Jordan Hill, Oxford OX2 8DP 200 Wheeler Road, Burlington MA 01803, Ontario, Canada
- Bunday BD, Mulholland H. 2014. *Pure Mathematics for Advanced Level* (Second edition). Elsevier
- Cooley HR. 2019. *Binomial Theorem*. McGraw Hill, USA
- Costa LI. 2017. A Linear Transformation to Accelerate the Convergence of the Negative Binomial Series. *Math. NA Winterthur*, Switzerland
- Dennis D, Addington S. 2009. The Binomial Series of Isaac Newton. *Mathematical Intentions*.
- Francia GA. 2000. Derivation and visualization of the binomial theorem. *International Journal of Computers for Mathematical Learning*, 1-24
- Gavrikov VL. 2018. Some Properties of Binomial Coefficients and their Application to Growth Modeling. *Arab Journal of Basic and Applied Sciences*, 25(1): 38-43
- Godman A, Telbert JF, Ogum GEO. 1984. *Additional Mathematics for West Africa*. Harlow, Longman, London, UK
- Goss D. 2011. *The Ongoing Binomial Revolution*. Number Theory. Springer
- Haggstrom P. 2012. *The Binomial Series for Negative Integral Exponents*. Maths at Bondi Beach, Sydney, Australia
- Horn ME. 2020. If you split something into two parts, you will get three pieces: The Bilateral Binomial Theorem and its consequences. *Journal of Physics: Conference Series*, 1612: 1-23
- Osanyinpeju KL, Aderinlewo AA, Dairo OU, Adetunji OR, Emmanuel SA, Ajisegiri ESA. 2019a. Development, Conversion and Application of Osanyinpeju (Power of Base 2) And Antiosanyinpeju (Antipower of Base 2) With Lekan (Power of Base 5) and Antilekan (Antipower of Base 5) Tables. 1st

- International Conference on Engineering and Environmental Sciences (ICEES) Theme: Advancing Technology and Environment Burdens (Challenges and Sustainable Solution. 969-981
- Osanyinpeju KL. 2019b. Development of Kifilideen (Power of Base 11) and Antikifilideen (Antipower of Base 11) Tables. 1st International Conference on Engineering and Environmental Sciences (ICEES) Theme: Advancing Technology and Environment Burdens (Challenges and Sustainable Solution). 957-968
- Osanyinpeju KL. 2020a. Development of Kifilideen Trinomial Theorem using Matrix Approach. International Conference on Electrical Engineering Applications (ICEEA 2020). 119-126, Department of Electrical Engineering, Ahmadu Bello University, Zaria, Kaduna State, Nigeria
- Osanyinpeju KL. 2020b. Utilization of Kifilideen Trinomial based on Matrix Approach in Computation of the Power of Base of Tri-Digits Number. International Conference on Electrical Engineering Applications (ICEEA 2020). 192-199, Department of Electrical Engineering, Ahmadu Bello University, Zaria, Kaduna State, Nigeria
- Osanyinpeju KL. 2020c. Computation of the power of base of two digits number using Kifilideen (Matrix, Combination and Distributive (MCD)) approach. Matrix Science Mathematics, 4(1): 14-19
- Osanyinpeju KL. 2020d. Development of Kifilideen (Power of Base 11), Antikifilideen (Antipower of Base 11) and Other Tables. Eliva Press SRL Publisher, MD-2060, bd. Cuza-Voda, ¼, of. 21 Chisinau, Republica Moldova
- Osanyinpeju KL. 2021a. Inauguration of Negative Power of $-n$ of Kifilideen Trinomial Theorem using Standardized and Matrix Methods. 2nd International Conference on Engineering and Environmental Sciences (ICEES) Theme: Re-thinking Engineering and Technology for Engineering Sustainability in the Face of Global Pandemic
- Osanyinpeju KL. 2021b. Inauguration of negative power of $-n$ of Kifilideen trinomial theorem using standardized and matrix methods. Acta Electronica Malaysia
- Osanyinpeju KL. 2021c. Establishment of Kifilideen Coefficient Tables for Positive and Negative of n and $-n$ of Kifilideen Trinomial Theorem Based on Matrix and Standardized Methods. Federal University of Agriculture, Abeokuta. Ogun State, Nigeria
- Schwartz RK. 2015. Newton's Binomial Theorem Marks 350 Years. Newsletter of the Schoolcraft College Mathematics Department, 23(3): 1-15
- Stroud KA, Booth DJ. 2007. Engineering Mathematics (6th Edition). Palgrave, London, UK
- Tavora M. 2020. The Unique Genius of Isaac Newton: A Beautiful Proof of a Well-Known Mathematical Result. Towards Data Science. <https://towardsdatascience.com>
- Telbert JF, Heng HH, John F. 1995. Additional Mathematics: Pure and Applied (6th Edition). Pearson Education South Asia, Philippines
- Tuttuh-Adegun MR, Adegoke MG. 2014. New Further Mathematics Project 2. Bounty Press Limited, Odutola Orita Challenge Police Station, Mapo, Ibadan, Nigeria
- Yang X. 2017. Engineering Mathematics with Examples and Applications: Binomial Theorem and Expansions. 31-35, Middlesex University, School of Science and Technology, London, UK
- Youngmee K, Sangwook R. 2014. The origin of Newton's Generalized Binomial Theorem. Journal of Mathematics, 27(2): 127-138