

Chronic Toxicological Evaluation and Reversibility Studies of *Moringa Oleifera* Ethanolic Seed Extract in Wistar Rats

¹Anoka A Njan, ²Olubunmi Atolani, ¹Olufunke E. Olorundare, ¹Saheed O Afolabi, ¹Benjamin C Ejimkonye, ¹Pender G Crucifix, ¹Babatunde Salami, ³Joseph O Olajide, ⁴Adeoye O Oyewopo and ¹Sabitiu A Oyeleke

¹Department of Pharmacology and Therapeutics, Faculty of Basic Medical Sciences, College of Health Sciences, University of Ilorin, Ilorin, Nigeria.

²Department of Chemistry, Faculty of Physical Sciences, University of Ilorin, Ilorin, Nigeria ³Department of Anatomy, Faculty of Basic Medical Sciences University of Ilorin, Ilorin Nigeria. ⁴Endocrinology Unit, Department of Anatomy Faculty of Basic Medical Sciences University of Ilorin Ilorin Nigeria

Abstract

This study evaluated the chronic toxicity and reversibility Potential of *Moringa oleifera* seed in Rats. Forty-four Wistar rats were randomized into four groups of 11 rats each, three groups were treated with doses of 50, 200, and 800 mg/kg of MOESE *p.o.* for 90 days while one group served as control and was administered distilled water *p.o.* After 90 days, half of the animals were sacrificed and blood samples collected for determination of biochemical and haematological parameters; antioxidants and *malondialdehyde* (MDA); sperm count, motility and morphology. Organs were harvested for weight determination and histopathological assessments the liver. The other half were left for another 30 days without treatment with the extract, after which they were sacrificed. MOESE produced significant reduction in food and water intake, relative weights of liver, heart, testes, lung and spleen at all treatment dose, these changes were overall small magnitude, within the range of historical control. There was significant increase in AST (200 and 800 mg/kg), LDH (200 mg/kg), ALP (800 mg/k), albumin (800 mg/kg) and K⁺ (800 mg/kg), but decrease in serum Ca²⁺ (50 and 200 mg/kg). A significant increase in serum antioxidant was observed at all treatment doses, while semen pH were reduced. These effects on organ relative weights and all biochemical parameters except AST and ALP were reversed after 30 days of the reversibility study. Histopathological presentations showed necrosis in the liver (50 and 200 mg/kg). Our study demonstrated that ethanol extract of *Moringa oleifera* seeds could cause infertility in male rats due to decreased semen pH. Our results also showed that the extract may be hepatotoxic due to persistent high level of AST and ALP even after cessation of exposure to the extract, this is also consistent with the histopathological results that shows hepatic necrosis associated with chronic exposure to the extract.

Key Words : *Moringa oleifera* seed, Toxicity, Wistar Rats

Introduction

Many plants have shown very effective medicinal values for various diseases in humans. As a result, pharmacological evaluation of medicinal plants has recently witnessed a growing interest amongst researchers worldwide. Studies have shown that several of the pharmaceuticals currently available have had a long history of use as herbal remedies.¹ Quinine, antimalarial drug from cinchona bark;² digoxin and digitoxin drugs for heart failure and atrial dysrhythmias from digitalis leaves^{1,3} are examples of the contributions of traditional pharmacopoeia. In Nigeria, despite immense progress in the delivery of health care at the primary health care level, an estimated 75 percent of the population still prefers to solve their health problems by consulting the traditional healers.⁴ In addition, herbal medicines are extensively used in the developing world where they offer a more affordable alternative to pharmaceutical drugs.

Moringa oleifera belongs to the family of moringaceae, a single genus with 14 known species.⁵ It is a fast growing drought resistant tree native of sub-Himalayan tracts of Northern India but now distributed worldwide in the tropics and sub tropics^{6,7} including Nigeria. The seed extracts of *Moringa oleifera* in particular have been reported to possess anti-inflammatory, antioxidant, anticancer, antimicrobial, Hypolipidaemic, antiulcer, antidiabetic, diuretic and hepatoprotective activities.⁸⁻¹³

Various studies have been conducted to investigate biological activities and efficacy of *Moringa oleifera* seed extracts,^{14,15} while little attention is being paid to its toxicity studies. Most of the studies on toxicity of *Moringa Oleifera* were focused on acute and sub-chronic toxicity¹⁴ and none of the studies has a report on reversibility of any toxic effect observed, from MOESE. This study therefore aims to investigate the chronic toxicity of MOESE and possibility of reversal of any toxic effect produced after 30 days of none treatment

Correspondence to:

Dr Anoka A Njan

Department of Pharmacology and Therapeutics,
Faculty of Basic Medical Sciences,
College of Health Sciences,
University of Ilorin, Ilorin, Nigeria.
Email: dinachim@yahoo.com
Telephone: 07033914915

Materials and methods

2.1 Identification and Preparation of Seeds Extract:
Moringa oleifera seed was bought from Avuco farms company Ltd, in Kaduna State Northwest Nigeria in




Fig.1: Effect of MOESE on body weight change of male rats in the main (90 day) study. Values are mean $\pm$ SEM. (n = 8).




Fig. 2: Effect of MOESE on body weight change of female rats in the main (90 day) study. Values are mean $\pm$ SEM. (n = 3).




Fig. 4: Effect of MOESE on daily water intake of rats in the main (90 day) study. Values are mean $\pm$ SEM. (n = 11).




Fig. 3: Effect of MOESE on daily food intake of rats in the main (90 day) study. Values are mean $\pm$ SEM. (n = 11).




Figure 5. Photomicrograph of transverse section of the liver of rats administered v distilled water and ethanol extracts of *M. oleifera* seed for 90 days. A: Distilled water treated liver shows normal hepatic plates and hepatocytes with normal histological features and intact stroma; B: 50 mg/kg body weight ethanol extract of *M. oleifera* treated liver shows dilated blood vessels and areas of marked hepatic necrosis; C: 200 mg/kg body weight ethanol extract of *M. oleifera* treated liver shows dilated blood vessels, sinusoidal haemorrhage and extensive areas of necrosis; D: 800 mg/kg body weight ethanol extract of *M. oleifera* treated liver shows dilated blood vessels and extensive areas of necrosis. H. magnification 400 $\times$.

Table 1: Effect of MOESE on relative organ weight of rats in the main (90 day) study

Dose (mg/kg)	Liver	Kidney	Lungs	Heart	Testes	Spleen	Pancreas	Brain
Control	3.01±0.02	0.29±0.01	0.79±0.02	0.32±0.00	0.74±0.08	0.39±0.01	0.25±0.02	0.68±0.03
50	2.66±0.06 ^a	0.28±0.01	0.71±0.02 ^a	0.29±0.01	0.82±0.01	0.29±0.01 ^a	0.28±0.02	0.70±0.01
200	2.53 ±0.02 ^{ab}	0.27±0.00	0.57±0.01 ^{ab}	0.27±0.01 ^a	0.48±0.01 ^{ab}	0.31±0.01 ^a	0.36±0.01 ^{ab}	0.77±0.04 ^a
800	2.79±0.01 ^{abc}	0.29±0.01	0.59±0.02 ^{ab}	0.31±0.01 ^c	0.49±0.02 ^{ab}	0.27±0.02 ^a	0.29±0.01 ^c	0.65±0.01 ^c

Values are mean ± SEM. (n = 5).

Table 2: Effect of MOESE on organ weight of rats in the reversibility study

Dose (mg/kg)	Liver	Kidney	Lungs	Heart	Testes	Spleen	Pancreas	Brain
Control	2.85±0.03	0.24±0.01	0.84±0.05	0.31±0.00	0.98±0.05	0.33±0.02	0.37±0.06	0.74±0.03
50	2.86±0.06	0.28±0.01 ^a	0.76±0.02	0.30±0.00	0.94±0.04	0.33±0.00	0.44±0.03	0.88±0.08
200	2.76±0.04	0.26±0.00 ^a	0.79±0.02	0.33±0.01	0.99±0.04	0.20±0.10	0.32±0.03	0.96±0.02 ^a
800	2.77±0.11	0.29±0.00 ^a	1.20±0.31	0.30±0.01	1.04±0.03	0.31±0.01	0.29±0.01	0.86±0.02

Values are mean ± SEM. (n = 5).

Table 3: Effect of MOESE on liver function parameters in rats in the main (90 day) study

Dose (mg/kg)	AST (u/L)	ALT (u/L)	ALP (u/L)	Protein (g/L)	Albumin (g/L)	TB (μmol/l)	CB (μmol/L)
0	234.33	47.33	84.67	47.33	18.33	8.33	0.23
	±1.20	±1.20	±1.76	±0.67	±0.33	±0.33	±0.03
50	223.00	46.00	84.33	52.00	20.67	8.33	0.367
	±21.00	±2.65	±0.88	±1.53	±1.20	±0.67	±0.03
200	356.33	45.00	81.33	48.00	20.33	9.00	0.33
	±22.66 ^{ab}	±2.65	±1.33	±0.58	±0.33	±0.58	±0.03
800	330.00	47.33	91.00	52.33	21.67	8.33	0.23
	±2.89 ^{ab}	±0.67	±0.58 ^{abc}	±3.18	±1.45 ^a	±0.33	±0.07

Values are mean ± SEM. (n = 5).

Table 4: Effect of MOESE on liver function parameters in rats in the reversibility study

Dose (mg/kg)	AST (u/L)	ALT (u/L)	ALP (u/L)	Protein (g/L)	Albumin (g/L)	TB (μmol/l)	CB (μmol/L)
0	139.67	42.33	90.33	66.67	23.00	0.57	0.20
	±0.88	±2.19	±1.33	±1.45	±0.00	±0.03	±0.05
50	137.33	37.00	92.00	69.00	25.67	0.53	0.23
	±0.33	±0.00	±1.00	±2.08	±0.88 ^a	±0.08	±0.03
200	268.67	41.00	90.33	65.33	24.00	0.57	0.23
	±0.67 ^{ab}	±0.58	±1.45	±2.85	±1.00	±0.03	±0.03
800	282.67	39.67	99.00	61.33	25.00	0.73	0.23
	±2.03 ^{abc}	±2.40	±0.58 ^{abc}	±1.76 ^b	±0.00	±0.12	±0.03

Values are mean ± SEM. (n = 5).

Table 5: Effect of MOESE on serum kidney function parameters in rats in the main (90 day) study

Dose (mg/kg)	Creatinine (μmol/L)	Urea (mmol/L)	Glucose (mmol/L)	Uric acid (mmol/L)
0	46.33 ± 1.45	8.30 ± 0.15	4.20 ± 0.15	0.38 ± 0.03
50	49.33 ± 2.03	8.23 ± 0.12	4.13 ± 0.15	0.44 ± 0.04
200	48.33 ± 2.40	8.47 ± 0.09	4.30 ± 0.15	0.51 ± 0.04
800	48.33 ± 1.67	8.47 ± 0.07	4.67 ± 0.39	0.43 ± 0.06

Values are mean ± SEM. (n = 5).

Table 6: Effect of MOESE on serum electrolytes in rats in the main study

Dose (mg/kg)	Na (mmol/L)	K (mmol/L)	Cl ⁻ (mmol/L)	HCO ₃ ⁻ (mmol/L)	Ca (mmol/L)
0	133.67± 1.20	6.60± 0.15	88.67± 1.45	21.00± 0.58	3.14± 0.05
50	134.00± 2.31	6.97± 0.12	89.67± 2.60	19.00± 1.52	2.72± 0.19 ^a
200	133.67± 0.33	6.77 ± 0.30	95.00± 1.00	20.33± 0.33	2.67± 0.10 ^a
800	135.00± 1.15	7.30 ± 0.23 ^a	95.00± 3.21	19.33± 0.88	2.82± 0.08

Values are mean ± SEM. (n = 5).

Table 7: Effect of MOESE on serum electrolytes in rats in the reversibility study

Dose (mg/kg)	Na (mmol/L)	K (mmol/L)	Cl ⁻ (mmol/L)	HCO ₃ ⁻ (mmol/L)	Ca (mmol/L)
0	135.00± 0.58	6.03± 0.83	98.67± 1.86	17.33± 0.33	2.68± 0.04
50	135.33± 1.20	6.97± 0.57	97.33± 0.88	18.67± 0.67	2.59± 0.09
200	135.00± 1.53	7.47± 0.30	89.67± 1.45 ^{ab}	22.00± 0.58 ^{ab}	2.74± 0.05
800	136.00± 1.73	7.10± 0.84	95.33± 1.76 ^c	20.00± 0.58 ^{ac}	2.67± 0.06

Values are mean ± SEM. (n = 5).

Table 8: Effect of MOESE serum lipid profiles of rats in the main study

Dose (mg/kg)	Trig (mmol/L)	Total chol (mmol/L)	LDL (mmol/L)	HDL (mmol/L)
0	1.13 ± 0.07	3.97 ± 0.03	1.01 ± 0.07	2.60 ± 0.15
50	1.17 ± 0.03	4.00 ± 0.10	0.96 ± 0.05	2.73 ± 0.15
200	1.20 ± 0.11	4.23 ± 0.09 ^{ab}	0.94 ± 0.12	3.17 ± 0.09 ^a
800	1.17 ± 0.09	3.73 ± 0.03 ^{abc}	1.03 ± 0.04	2.60 ± 0.09 ^c

Values are mean ± SEM. (n = 5).

Table 9: Effect of MOESE on serum lipid profiles of rats in the reversibility study

Dose (mg/kg)	Trig (mmol/L)	Total chol (mmol/L)	LDL (mmol/L)	HDL (mmol/L)
0	0.61 ± 0.03	4.63 ± 0.32	2.50 ± 0.25	2.50 ± 0.25
50	0.68 ± 0.12	4.43 ± 0.07	2.87 ± 0.03	2.87 ± 0.03
200	0.80 ± 0.02	4.43 ± 0.15	2.77 ± 0.15	2.77 ± 0.15
800	0.71 ± 0.10	4.40 ± 0.10	2.17 ± 0.27	2.17 ± 0.27

Values are mean ± SEM. (n = 5).

Table 10: Effect of MOESE on antioxidants and MDA levels in serum (90 day study)

Dose (mg/kg)	SOD (μl/ml)	GPX (μl/L)	MDA (μmol/L)
0	208.33±0.29	2399.70±45.11	3.07±0.07
50	277.67±5.55 ^a	3025.30±43.56 ^a	2.43±0.15 ^a
200	229.33±4.84 ^b	2561.00±21.03 ^{ab}	2.13±0.15 ^a
800	210.00±7.64 ^b	3248.00±44.00 ^{abc}	2.30±0.30 ^a

Values are mean ± SEM (n=5).

Table 11: Effect of MOESE on antioxidants and MDA level in serum (30 day reversibility study)

Dose (mg/kg)	SOD (ul/ml)	GPX (ul/L)	MDA (umol/L)
0	204.33±5.81	2699.30±186.2	3.40±0.23
50	277.33±4.40 ^a	3083.00±19.76	1.97±0.03 ^a
200	309.00±3.06 ^{ab}	3407.30±196.36 ^a	2.03±0.03 ^a
800	256.00±8.00 ^{abc}	3186.00± 282.00	2.20±0.50 ^a

Table 12: Effect of MOESE on hematological parameters in blood (90 day study)

Dose (mg/kg)	PCV (%)	WBC (10 ⁹ /L)	N (%)	L (%)	RBC (10 ¹² /L)	Hb (g/dl)	PLT (10 ⁹ /L)	MCV (fl)	MCH (pg)	MCHC (g/dl)
0	36.67 ±0.33	5.51 ±0.47	69.00 ±0.57	30.33 ±0.33	6.26 ±0.06	15.77 ±0.22	255.00 ±24.70	59.00 ±0.00	26.43 ±0.13	43.93 ±0.68
50	36.67 ±0.33	6.87 ±0.79	69.67 ±0.33	30.00 ±0.33	6.27 ±0.11	15.83 ±0.37	359.67 ±25.34	56.33 ±0.88	25.53 ±0.18	44.73 ±1.14
200	35.00 ±1.15	4.95 ±0.38	69.67 ±0.33	30.00 ±0.00	6.24 ±0.08	17.20 ±0.17	312.00 ±50.14	55.33 ±0.33*	27.40 ±0.35	45.53 ±1.59
800	33.00 ±3.00	7.28 ±1.13	68.33 ±1.86	33.50 ±0.50	6.55 ±0.18	16.20 ±1.10	312.50 ±30.50	55.00 ±1.00*	27.00 ±0.20	48.95 ±0.65

Values are expressed as mean ± S.E.M. (n = 5).

Table 13: Effect of MOESE on some hematological parameters in blood (30 day reversibility study)

Dose (mg/kg)	PCV (%)	WBC (10 ⁹ /L)	N (%)	L (%)	RBC (10 ¹² /L)	Hb (g/dl)	PLT (10 ⁹ /L)	MCV (fl)	MCH(pg)	MCHC (g/dl)
0	45.00 ±0.58	10.90 ±1.39	68.66 ±0.67	31.33 ±0.67	8.59 ±0.06	15.77 ±0.33	412.67 ±35.41	63.33 ±1.45	22.37 ±0.34	36.27 ±0.74
50	43.67 ±1.33	10.63 ±1.78	68.33 ±1.20	30.67 ±0.33	6.86 ±0.18	15.80 ±0.71	385.67 ±36.73	63.67 ±0.66	22.97 ±0.48	36.07 ±0.64
200	45.00 ±1.53	6.10 ±1.15	85.33 ±0.33*	35.33 ±0.33*	4.97 ±0.09*	15.33 ±0.79	404.33 ±31.15	64.67 ±1.33	21.20 ±0.49	30.33 ±0.23
800	38.50 ±0.50	5.99 ±0.29	65.50 ±0.50	34.50 ±0.50*	5.99 ±0.41	11.55 ±0.45	377.50 ±20.50	64.50 ±3.50	23.10 ±1.50	38.75 ±3.35

Values are expressed as mean ± S.E.M. (n = 5).

Table 14: Effect of MOESE on sperm motility, count, morphology and pH (90 day study)

Dose (mg/kg)	Sperm motility (%)	Sperm count (millions/ml)	Abnormality (%)	pH
0	75.33±7.75	39.67±0.67	7.20±1.72	4.70±0.20
50	66.67±6.67	42.33±0.88	8.33±2.01	4.20±0.06*
200	75.00±0.00	36.00±1.15	7.40±1.50	4.17±0.03*
800	57.50±2.50	41.00±6.00	5.50±1.71	4.25±0.05*

Values are expressed

of 50mg/kg caused statistically significantly increase ($p < 0.05$) in body weight on day 35, compared to control while a statistically significant reduction ($p < 0.05$) in body weight at dose of 800mg/kg on day 49 were observed, compared to 50mg/kg. However, in the female rats, no statistically significant differences in body weight were observed at all doses.

The relative organ weights of the rats are shown in Table 1. At doses 50, 200, and 800 mg/kg, there were statistically significant reduction ($p < 0.05$) in the relative organ weights of the liver, lungs and spleen, compared to control. There was also statistically significant reduction ($p < 0.05$) in weight of testes at doses of 200 and 800 mg/kg, compared to control and 50mg/kg groups, while the relative weight of the heart, at the dose of 200 mg/kg, was statistically significantly reduced ($p < 0.05$), compared to control. There was a statistically significant increase ($p < 0.05$) in the relative weights of pancreas and brain at the dose of 200 mg/kg compared to control respectively. When compared to 50 mg/kg, there was a statistically

significant reduction ($p < 0.05$) in the relative weight of liver (at dose of 200 mg/kg), lungs and testes (at the doses of 200 and 800 mg/kg), while a statistically significant increase ($p < 0.05$) in the weights of liver (at the dose of 800 mg/kg) and pancreas (at the dose of 200 mg/kg) were also observed when compared to 50mg/kg.

When compared with 200 mg/kg, there was a statistically significant increase ($p < 0.05$) in weights of liver and heart, and significant reduction ($p < 0.05$) in the weights of pancreas and brain, at the dose of 800 mg/kg respectively. The significant differences observed were reversed after 30 days of cessation of treatment (Table 2).

Effect of MOESE on Food and Water Intake

Food consumption and water intake of rats treated with MOESE doses (50, 200 and 800) for 90 days are presented in Figures 3 and 4 respectively. In respect of food intake, compared to control, MOESE produced significant reduction ($p < 0.05$) in average

daily food intake at dose of 50 mg/kg on days 20, 30, 40, and 70. The same trend of effect (significant reduction) was obtained ($p < 0.05$) at doses of 200 mg/kg and 800 mg/kg but throughout the treatment period. Compared to 50 mg/kg treatment group, there was significant reduction ($p < 0.05$) in average daily food intake at doses of 200 mg/kg and 800 mg/kg all through the treatment period except on days 80 and 90 when there was no significant difference ($p > 0.05$) when 200 mg/kg was compared to 50 mg/kg. Compared to 200 mg/kg, dose of 800 mg/kg produced significant decrease ($p < 0.05$) in average daily food intake on days 10, 60 and 80.

Concerning water intake, compared to control, MOESE at the dose of 50 mg/kg produced significant reduction ($p < 0.05$) in average daily water intake on days 10, 20, 30, 40, 50, and 90. Also, there was significant reduction at doses of 200 mg/kg and 800 mg/kg throughout the treatment period (days 10, 20, 30, 40, 50, 60, 70, 80 and 90).

EFFECT OF MOESE ON BIOCHEMICAL PARAMETERS

Effect of MOESE on Liver Function Parameters

There was significant increase ($p < 0.05$) in the activity of aspartate aminotransferase (AST) in the groups treated with 200 mg/kg (356.33 ± 22.66 U/L) and 800 mg/kg (330 ± 2.89 U/L) when compared to control (234 ± 1.20 U/L) (Table 3). When compared to 50 mg/kg treatment group, there was significant increase ($p < 0.05$) in AST activity in the groups treated with 200 mg/kg and 800 mg/kg. There was also a significant increase ($p < 0.05$) in the activity of alkaline phosphatase (ALP) in the groups treated with 800 mg/kg (91.00 ± 0.58) when compared to control (84.67 ± 1.76). A similar trend of increase occurred when compared 800 mg/kg to 50 mg/kg (84.33 ± 0.88) and to 200 mg/kg (81.33 ± 1.33) treatment groups. The concentration of albumin was significantly increased ($p < 0.05$) in the group treated with *M. oleifera* seed at the dose of 800 mg/kg (21.67 ± 1.45) when compared to control (18.33 ± 0.33). The effects on AST and ALP were not reversed after 30 days of cessation of administration of the ethanol extract of *M. oleifera* seed because the significant increase observed after 90 days administration still persisted (Table 4). The effect on albumin was, however, reversed after 30 days of cessation of administration of the extract (Table 4).

Effect of MOESE on Kidney Function Parameters

There was a no significant difference ($p > 0.05$) in serum levels of these parameters at all treatment doses when compared to the control group (Table 5).

Effect of MOESE on Serum Electrolytes Concentrations

There was a significant increase ($p < 0.05$) in

concentration of potassium (K) in the group treated with 800mg/kg of the extract (7.30 ± 0.23 mmol/L) when compared to the control group (6.60 ± 0.15 mmol/L) (Table 6). There was significant reduction ($p < 0.05$) in calcium concentrations in the groups treated with 50 mg/kg and 200 mg/kg of the extract when compared to the control. The effects on potassium and calcium were reversed after 30 days of cessation of the extract administration, while chloride concentration was significantly reduced in 200 mg/kg treatment group and bicarbonate was significantly increased in 200 mg/kg and 800 mg/kg groups (Table 7).

Effect of MOESE on Lipid Profiles

There was a significant increase ($p < 0.05$) in the concentration of total cholesterol in the group treated with the extract at the dose of 200 mg/kg (4.23 ± 0.09 mmol/L) when compared to the control group (3.97 ± 0.03 mmol/L); the increment ($p < 0.05$) is also significant when compared to 50 mg/kg treatment group (4.00 ± 0.10 mmol/L) (Table 8). A significant reduction ($p < 0.05$) in concentration of total cholesterol was obtained in the group treated with 800 mg/kg (3.73 ± 0.03 mmol/L) when compared to the control group (3.97 ± 0.03 mmol/L) and to the 50 mg/kg (4.00 ± 0.10 mmol/L) and 200 mg/kg (4.23 ± 0.09 mmol/L) treatment groups (Table 8). There was also a significant increase ($p < 0.05$) in the concentration of HDL in the groups treated with 200 mg/kg (3.17 ± 0.09 mmol/L) of the extract when compared to the control group (2.60 ± 0.15 mmol/L). At the dose of 800 mg/kg, there was a significant decrease ($p < 0.05$) in HDL concentration when compared to 200 mg/kg but not significantly different from the control group (Table 8). The effects on total cholesterol and HDL were reversed after 30 days of cessation of administration of the extract (Table 9).

Effect of MOESE on Serum Antioxidants and Malondialdehyde Levels

At doses of 200mg/kg and 800mg/kg, there was no significant difference in the level of superoxide dismutase (SOD) in serum. However, a significant increase ($p < 0.05$) in the level of glutathione peroxidase (GPX) was observed with 200mg/kg dose of MOESE (2561.00 ± 21.03 ul/L vs control value of 2399.70 ± 45.11 ul/L) (Table 10). Also, there was a significant increase ($p < 0.05$) in GPX with the 800mg/kg dose of MOESE (3248.00 ± 44.00 vs control value of 2399.70 ± 45.71) (Table 10). Also, there was a significant decrease ($p < 0.05$) in the level of SOD in the group that received 200mg/kg dose of MOESE (229.33 ± 4.84 ul/ml) compared to the 50mg/kg dose of MOESE (277.67 ± 5.55 ul/ml) (Table 10). A significant decrease ($p < 0.05$) in the level of SOD in the group that received 800mg/kg dose of MOESE (210.00 ± 7.64 ul/ml) when compared to the group that received

50mg/kg dose of MOESE (277.67 ± 5.55 ul/ml) was also recorded (Table 10).

At the dose of 50mg/kg, there was a significant increase ($p < 0.05$) in the level of SOD (277.67 ± 5.55 vs control value of 208.33 ± 9.28 ul/ml), as well as a significant increase ($p < 0.05$) in the level of glutathione peroxidase (GPX), (3025.30 ± 43.56 ul/L vs control value of 2399.70 ± 45.71 ul/L) (Table 10). Also, there was a significant decrease ($p < 0.05$) in the level of GPX in the group that received 200mg/kg dose of MOESE (2561.00 ± 21.03 ul/L) when compared to the group that received 50mg/kg dose of MOESE (3025.30 ± 43.56 ul/L) (Table 10). There was a significant increase ($p < 0.05$) in the level of GPX in the group that received 800mg/kg dose of MOESE (3248.00 ± 44.00 ul/L) when compared to the group that received 50mg/kg dose of MOESE (3025.30 ± 43.56 ul/L) (Table 10). There was also a significant increase ($p < 0.05$) in the level of GPX in the group that received 800mg/kg dose of MOESE (3248.00 ± 44.00 ul/L) when compared to the group that received 200mg/kg dose of MOESE (2561.00 ± 21.03 ul/L) (Table 10).

There were significant decreases in the level of MDA across the treatment group after the 90 day period (Table 10). At the dose of 50mg/kg, MOESE caused a significant decrease ($p < 0.05$) in the level of MDA (2.43 ± 0.15 vs control value of 3.07 ± 0.07 umol/L) (Table 10). There was a significant decrease ($p < 0.05$) in the level of MDA in the group treated with MOESE at the dose of 200mg/kg (2.13 ± 0.15 vs 3.07 ± 0.07 umol/L) compared with the distilled water treated group (Table 10). A significant decrease ($p < 0.05$) in the level of MDA was also observed in the group treated with MOESE at the dose of 800mg/kg (2.30 ± 0.30 vs control value of 3.07 ± 0.07 umol/L) (Table 10).

After a 30 day cessation of treatment, there were significant increases in the level of superoxide dismutase (SOD) enzyme in the serum across the treatment groups (Table 11); this suggests that the increase in free radical generation mediated by MOESE were retained even with cessation of therapy. At the dose of 50mg/kg, there was a significant increase ($p < 0.05$) in the level of SOD (277.33 ± 4.40 vs 204.33 ± 5.81 ul/ml in control) (Table 11), at 200mg/kg, there was also a significant increase ($p < 0.05$) in the level of SOD (309.00 ± 3.06 vs 204.33 ± 5.81 ul/ml in control) (Table 11), and at 800mg/kg, a significant increase ($p < 0.05$) in the level of SOD (256.00 ± 8.00 vs 204.33 ± 5.81 ul/ml in control group) was also recorded (Table 11). There was a significant increase ($p < 0.05$) in the level of SOD in the group that earlier received 200mg/kg dose of MOESE (309.00 ± 3.06 ul/ml) when compared with the group that was previously treated with 50mg/kg dose of MOESE (277.33 ± 4.40 ul/ml) (Table 11). There was a significant decrease ($p < 0.05$) in the level of SOD in the group that previously

received 800mg/kg dose of MOESE (256.00 ± 8.00 ul/ml) when compared with the 50mg/kg dose of MOESE previously treated animals (277.33 ± 4.40 ul/ml) (Table 11). A significant decrease ($p < 0.05$) in the level of SOD was also recorded in the group that received 800mg/kg dose of MOESE (256.00 ± 8.00 ul/ml) when compared with the group that received 200mg/kg dose of MOESE (309.00 ± 3.06 ul/ml) (Table 11). The group treated with 200mg/kg dose of MOESE showed a significant increase ($p < 0.05$) in the level of GPX (3407.30 ± 196.36 vs 2699.30 ± 186.2 ul/L) when compared with the group treated with distilled water (Table 11). There were significant decreases in the level of MDA across the treated group after the 30 day cessation of therapy (Table 11). The group initially treated with MOESE at the dose of 50mg/kg showed a significant decrease ($p < 0.05$) in the level of MDA (1.97 ± 0.03 vs 3.40 ± 0.23 umol/L) when compared with the control group (Table 11). There was also a significant decrease ($p < 0.05$) in MDA level in the group that initially received 200mg/kg dose of MOESE (2.03 ± 0.03 vs control value of 3.40 ± 0.23 umol/L) (Table 11). A significant decrease ($p < 0.05$) in the level of MDA was also observed in the group that previously received 800mg/kg dose of MOESE (2.20 ± 0.50 vs control value of 3.40 ± 0.23 umol/L) after the 30 day cessation of therapy (Table 11).

Effect of MOESE on Haematological Parameters in Blood

MOESE did not produce any significant effect ($P > 0.05$) on haematological parameters after the 90 day administration except for MCV in which there was a significant decrease ($p < 0.05$) in the group treated with MOESE at the dose of 200mg/kg (55.33 ± 0.33 10^9 /L) when compared with the group treated with distilled water (59.00 ± 0.00 10^9 /L) (Table 12), and also a significant decrease ($p < 0.05$) in the group treated with 800mg/kg *M. oleifera* (55.00 ± 1.00 vs control value of 59.00 ± 0.00 10^9 /L) (Table 12).

After the 30 day cessation of treatment, a significant increase ($p < 0.05$) was recorded in the level of neutrophils in the group earlier treated with 200mg/kg dose of MOESE ($85.33 \pm 0.33\%$) when compared with the distilled water treated group (68.66 ± 0.67) (Table 13). Also, a significant increase ($p < 0.05$) in the level of lymphocytes in the group earlier treated with 200mg/kg dose of MOESE (35.33 ± 0.33 vs $31.31 \pm 0.67\%$ in control group) was recorded (Table 13). A significant increase ($p < 0.05$) in the level of lymphocytes was also recorded in the group that earlier received 800mg/kg dose of MOESE (34.50 ± 0.50 vs control value of $31.33 \pm 0.67\%$) (Table 13). For RBC, a significant decrease ($p < 0.05$) in the group earlier treated with 200mg/kg dose of MOESE (4.97 ± 0.09 10^{12} /L) compared with control value of (5.59 ± 0.06 10^{12} /L), was also observed (Table 13).

EFFECT OF MOESE ON SPERM MOTILITY, COUNT, MORPHOLOGY AND pH

MOESE did not produce any significant effect on sperm motility, count and abnormality after the 90 day administration period. However, there was significant decrease in pH across all the treated groups compared to control animals (Table 14). At the dose of 50mg/kg of MOESE, there was a significant decrease ($p < 0.05$) in pH (4.20 ± 0.06 vs control value of 4.70 ± 0.20) (Table 14). A significant decrease ($p < 0.05$) in pH was also observed in the group treated with 200mg/kg dose of MOESE (4.17 ± 0.03) when compared with the distilled water treated group (4.7 ± 0.20) (Table 14). It was also observed that the 800mg/kg dose of MOESE treated group showed a significant decrease ($p < 0.05$) in pH (4.25 ± 0.05) when compared with the distilled water treated group (Table 14).

After the 30 day cessation of therapy, no significant difference in sperm parameters was observed (Table 15).

Effect of *M. oleifera* seed ethanolic extract on histological presentations

Concerning the liver, there was no adverse histopathological presentations observed in the control group (Fig. 5A). The liver appeared normal with preserved hepatic architecture, hepatocytes with normal histological features and intact stroma. The liver of rats that received 50 mg/kg presented with dilated blood vessels, areas of marked necrosis and hyperchromatic foci (Fig. 5B), while that of 200 mg/kg showed dilated blood vessels, sinusoidal haemorrhage and extensive areas of necrosis (Fig. 5C). Dilated blood vessels and extensive areas of necrosis were also observed in the liver at the doses of 800 mg/kg (Fig. 5D).

Discussion

The reticence to the widespread use of medicinal plants is gradually being eroded by the inability of orthodox drugs to cure long standing and common ailments such as diabetes and hypertension. Additionally, anecdotal evidence of efficacy of medicinal plants has boosted confidence in the use of such products. In the present study, the 96h LD₅₀ value of *M. oleifera* seed extract for the rats was found to be 2900mg/kg which indicates that *M. oleifera* seed extract could be relatively not toxic to rats at this concentration based on chemical labeling and clarification of acute systemic toxicity on oral LD50 values recommended by the Organization for Economic Cooperation and Development (OECD).³¹ However, it has been reported that the LD50 is not an absolute value but an inherently variable biologic parameter that cannot be compared to constants sure as

molecular weight or melting point.³² Organ weight changes can be sensitive indicators of target organ toxicity, and significant changes in organ weights may occur in the absence changes in other pathology parameters.³³ In this study, there were no body weight, body weight gain, food consumption, absolute or relative organ weight, or water consumption or clinical changes associated with the administration of MOSES. Any body weight and food and water consumption changes were overall small magnitude, within the range of historical control (Table 1&2, Fig 1, 2&3). Liver function tests help in the diagnosis of any abnormal/normal condition of liver. Leakage of cellular enzymes into plasma indicates the sign of hepatic tissue damage.^{34,35} Generally measurement of alanine aminotransferase (ALT) and AST are used as important diagnostic marker to indicate liver injury due to hepatotoxins. In this study, administration of MOSES extract significantly ($p < 0.05$) increase serum AST, ALP and albumin levels but did not changed ALT values in treated animals relative to control (Table 3 & 4). The histology results also shows the damage of parenchymal and mitochondrial sites of the liver where ALT is located. Our results is dissimilar to that of Ferreira et al (2009)³⁶ who reported no alterations in hepatic enzymes in rat treated with 1300-1670mg/kg/day *Moringa Oleifera* seed for 30days, but agrees with Bharani et al. (2003)³⁷ who reported that administration of the hydroalcoholic extract of *M. oleifera* drumsticks by oral route elevated levels of some hepatic enzymes involved in detoxification of xenobiotic substances, such as carcinogens and plant venomous compounds. The increased serum albumin level suggests a potential for the seed to cause imbalance fluid distribution. Albumin plays a major role in stabilizing extracellular fluid volume by contributing to oncotic pressure of plasma. Because smaller animals (for example rats) function at a lower blood pressure, they need less oncotic pressure to balance this, and thus need fewer albumins to maintain proper fluid distribution.³⁸ The increment in AST and ALT persisted while that of albumin was reversed after 30 days of cessation of the extract.

Reactive oxygen species (ROS) are related to oxidative stress and many scientific reports have shown that excessive production of ROS can further aggravate oxidative stress and have implicated ROS in a number of disorder/disease processes, including heart disease, diabetes, liver injury, cancer, and aging.³⁹⁻⁴¹ Equilibrium between ROS and enzymatic antioxidant enzymes including superoxide dismutase (SOD) and glutathione peroxidase (GPx) are crucial and could be an important mechanism for preventing damage by oxidative stress. The liver plays a central role in the maintenance of systemic lipid homeostasis and is especially susceptible to ROS damage, mechanism has been suggested to play role in preventing APAP toxicity.⁴² In

this study, MOSES has shown a significant increase of these enzymes (Tables 10 & 11) suggesting the presence of antioxidants compounds. These results is consistent with previous studies described by Gupta et al. (2005)⁴³ and his colleagues that shows the capacity of *moringa* seeds to protect animals against oxidant stress caused by arsenic exposure. Several researches attributed this oxidative protection of the seeds to the presence of compounds with antioxidant activity against free radicals.^{44,45}

No significant effect was produced by MOESE on hematologic parameters after the 90 day administration except a slight decrease in MCV in the group treated with MOESE at the dose of 200mg/kg and 800mg/kg when compared with the control group, the decrease in MCV was however reversed after the 30 day cessation of treatment. (Tables 12 & 13)

Regarding the semen analysis, significant differences were not observed in sperm count, motility and abnormality across the three treatment doses of MOESE used during the 90 day treatment period when compared with control animals. However, there was a significant decrease in the semen pH in all treated group when compared to control group (Table 14). The microenvironment of sperm, seminal plasma, is of great significance. Seminal plasma is a mixture of secretions from the testes, epididymides and accessory sex glands. Seminal plasma contains $\text{HCO}_3^-/\text{CO}_2$, inorganic ions, organic acids, sugars, lipids, steroids, amino acids, polyamines, nitrogenous bases and proteins.⁴⁶ As a result, semen has a very high buffering capacity, higher than that of most other body fluids. Therefore, the pH of the seminal fluid may play a significant role not only in maintaining the viability and quality of the sperm, but also in ensuring fertilization. Declined Na^+/K^+ -ATPase activity at acidic pHs has been reported to result in decreased sperm movement and capacitation, which could be one of the mechanisms of male infertility,⁴⁷ this study shows marked decrease in the pH of sperm in animals treated with MOSES when compared with control, suggesting that the extract at given doses may cause infertility in male rats. The decrease in pH was reversed after the 30 day cessation of treatment in all three treated groups. Histopathology of the liver from treated rats showed degenerative lesions on hepatic tissue. Some hepatotoxic drugs produce their effects through toxic metabolites and immunological reaction,⁴⁸ This result is buttressed by increased level of AST and ALP.

Our study clearly demonstrated that ethanol extract of *Moringa oleifera* seeds could cause infertility in male rats due to decreased semen pH. Our results also showed that the extract may be hepatotoxic due to persistent high level of AST and ALP even after cessation of exposure to the extract, this is also consistent with the histopathological results that shows

hepatic necrosis associated with chronic exposure to the extract.

Acknowledgements

This work was supported by grant from Raw Material Research and Development Council (RMRDC) Abuja, Nigeria. The authors are grateful to Late Professor F.A Oladele for authenticating the plant material, Mr. Adesua Alfred Obafemi and Dr. Haas Akanbi of Department of Pharmacology and Therapeutics, Faculty of Basic Medical Sciences, University of Ilorin, Ilorin, Kwara State, Nigeria for their remarkable assistance during the course of this study.

References

1. Kebede S, Afework M, Debella A, Ergete M, Makonnen E. Toxicological study of the b u t a n o l fractionated root extract of *Asparagus africanus* Lam., on some blood parameter and histopathology of liver and kidney in mice. *BMC Research Notes*. 2016; 9(49):1-15.
2. Himanshu M, Bhupendra KM, Dharam CJ. Optimization of extraction conditions and hptlc - uv method for determination of quinine in different extracts of *Cinchona species* bark. *Rec Nat Prod*. 2008; 2(4): 107-115.
3. Smith TW, Antman EM, Friedman PL, Blatt CM, Marsh JD. Digitalis glycosides: mechanisms and manifestations of toxicity. *Prog Cardiovasc Dis*. 1984; 26:413- 458.
4. Adefolaju T. (2011). The dynamics and challenging structure of traditional healing system in Nigeria. *International Journal of Health Research*. 2011; 4(2):99-106.
5. Fahey H. *Moringa oleifera*: A Review of the Medical Evidence for Its Nutritional, Therapeutic, and Prophylactic Properties. Part 1. *TreesLife J*. 2005; 1:5.
6. Njan AA, Amali MO, Olatunji LO, Olorundare OE. An Overview of the Ethno- Pharmacological Potentials of *Moringa Oleifera* Lam, "The Miracle Tree". *Archives of Basic & Applied Medicine*. 2014; 2:135-145.
7. Aja PM, Ibiam UA, Nwali BU, Orji OU, Edwin N, Afiukwa CA. Evaluation of Effect of E t h a n o l Extracts of *Moringa oleifera* and *Cajanus cajan* Leaves on Lipid Profiles in Alloxan Induced Diabetic Albino Rats. *Global Journal of Biotechnology & Biochemistry*. 2015; 10(2):71-76.
8. Gupta R, Mathur M, Bajaj VK, Katariya P, Yadav S, Kamal R, Gupta RS. Evaluation of antidiabetic and antioxidant activity of *Moringa oleifera* in experimental diabetes. *Journal of Diabetes*. 2012; 4:164-171.

9. Sharida Fakurazi, Syazana Akmal Sharifudin, Palanisamy Arulselvan. Moringa oleifera Hydroethanolic Extracts Effectively Alleviate Acetaminophen-Induced Hepatotoxicity in Experimental Rats through Their Antioxidant *Nature Molecules* 2012; 17:8335-8350.
10. Alikwe PCN, Ohimain EI, Zige DV, Angaye TNC (2013). Antibacterial activity of ethanol extract of the defatted seed and seed coat of Moringa oleifera. *IOSR Journal of Pharmacy and Biological Sciences* 2013; 8(1):38-41.
11. Kansara SS, Singhal M. Evaluation of Antiulcer Activity of Moringa Oleifera Seed Extract. *Journal of Pharmaceutical Science and Bioscientific Research* 2013; 3(1):20-25.
12. Minaiyan M, Asghari G, Taheri D, Saeidi M, Nasr-Esfahani S. Anti-inflammatory effect of Moringa oleifera Lam. seeds on acetic acid-induced acute colitis in rats. *Avicenna Journal of Phytomedicine*. 2014; 4(2):127-136.
13. Atsukwei D, Eze ED, Moses DM, Adinoyi SS, Upkabi CN. Hypolipidaemic effect of ethanol leaf extract of Moringa Oleifera Lam. in experimentally induced hypercholesterolemic wistar rats. *Int. J. Nutri. And Food Sci.* 2014; 3(4):355-360.
14. Ajibade TO, Arowolo R, Olayemi FO. Phytochemical screening and toxicity studies on the methanol extract of the seeds of Moringa oleifera. *J. Complement Integr. Med.* 2013; 10:11-16.
15. Sana A, Saleem R, Faizi S. Hypotensive Activity of Moringa oleifera Lam (Moringaceae) Root Extracts and its Volatile Constituents. *Tropical Journal of Pharmaceutical Research*. 2015; 14(5):823-830.
16. Harbone JB. Phytochemical Methods London. Chapman and Hall Ltd. 1973; 49-188.
17. Bargah RK. Preliminary test of phytochemical screening of crude ethanolic and aqueous extract of Moringa pterygosperma Gaertn. *Journal of Pharmacognosy and Phytochemistry*. 2015; 4(1):07-09
18. Guide for the Care and Use of Laboratory Animals. Revised. NIH publication. 1978; No. 83-23. National Institute of Health Bethesda. MD.
19. Lorke D. (1983). A new approach to practical acute toxicity testing. *Arch. Toxicol.* 1983; 54(4):275-278
20. Afolabi SO, Akindele AJ, Awodele O, Anunobi CC, Adeyemi OO. A 90 day chronic toxicity study of Nigerian herbal preparation DAS-77 in rats. *BMC Complementary and Alternative Medicine* 2012; 12:79.
21. Gornall AC, Bardawill CJ, David MM. Determination of serum protein by means of biuret reaction. *J Biol Chem.* 1949; 177(2):751-756.
22. Plummer DT. An introduction to practical biochemistry. McGraw-Hill. 1978; 119-154.
23. Baginsky ES, Marie SS, Clark WL, Zak B. Direct microdetermination of calcium. *Clinica Chimica Acta.* 1973; 46:49-54.
24. Cheesbrough M. District Laboratory Practice in Tropical Countries, Part 1. UK: Cambridge University Press. 2005; 365-370.
25. Ikewuchi JC, Onyeike EN, Uwakwe AA, Ikewuchi CC. Effect of aqueous extract of the leaves of Acalypha wilkesiana 'Godseffiana' Muell Arg (Euphorbiaceae) on the hematology, plasma biochemistry and ocular indices of oxidative stress in alloxan induced diabetic rats. *J Ethnopharmacol* 2011; 137:1415-1424.
26. Soon YY, Tan BKH. Evaluation of the hypoglycemic and antioxidant activities of Morinda officinalis in streptozocin-induced diabetic rats. *Singapore Med J.* 2002; 43:077-085.
27. Cheesbrough M. Examination of semen. In District laboratory practice in tropical countries (Part 2). UK: Cambridge University Press. 2000; 130-132
28. Ogli SA, Enyikwola O, Odeh SO. Evaluation of the efficacy of separate oral supplements compared with the combined oral supplements of vitamins C and E on sperm motility in Wistar rats. *Niger J Physiol Sci* 2009; 24:129-135.
29. Krause WJ. The art of examining and interpreting histologic preparations. A student handbook. UK: Parthenon Publishing Group. 2001; 9-10.
30. Mahajan BK. Significance of Difference in Means. In: Mahajan BK, editor. Methods in Biostatistics for Medical and Research Workers. JAYPEE Brothers Medical Publishers. 1997; 130-55.
31. Walum, E. Acute oral toxicity. *Environ. Health Perspect.* 1998; 106:497-502.
32. Oliver JA. Opportunity for using fewer animals in acute toxicity studies. In: *Chemicals Testing and Animal Welfare*. The National Chemicals Inspectorate, Solana, Sweden. 1986; 119-143.
33. Bailey SA, Zidell RH, Perry RW. Relationship between organ weight and bodyweight/brain weight in the rat: what is the best analytical endpoint? *Toxicol. Pathol.*, 2004; 32:448.
34. Ramaiah SK. A toxicologist guide to the diagnostic interpretation of hepatic biochemical parameters. *Food Chem. Toxicol.* 2007; 45:1551-1557.
35. Sreelatha S, Padma PR, Umadevi M. Protective effects of Coriandrum sativum extracts on carbon tetrachloride-induced

- hepatotoxicity in rats. *Food Chem. Toxicol.* 2009; 47 : 702–708.
36. Ferreira PMP, Carvalho AFU, Farias DF, Cariolano NG, Melo VMN, Quiroz MGR, Martins AMCM, Machado-Neto JC. Larvicidal activity of the water extract of moringa oleifera seeds against Aedes aegypti and its toxicity upon laboratory animals. *An Acad. Bras. Ciencias.* 2009;81(2):207-216.
 37. Bharali R, Tabassum J, and Azad M.R. Chemomodulatory effect of Moringa oleifera Lam. on hepatic carcinogen metabolizing enzymes, antioxidant parameters and skin papillomagenesis in mice. *Asian Pac J Cancer Prev* 2003; 4 :131-139.
 38. Harper ME, Dugaiczyk A. *Am. J. Hum. Genet.* 1983; 35(4):565–572.
 39. Sies H, Stahl W, Sevanian A. Nutritional, dietary and postprandial oxidative stress. *J. Nutr.* 2005; 135:969–972.
 40. Giordano FJ. Oxygen, oxidative stress, hypoxia, and heart failure. *J. Clin. Invest.*, 2005; 115:500–508.
 41. Klaunig JE, Kamendulis LM. The role of oxidative stress in carcinogenesis. *Annu. Rev. Pharmacol. Toxicol.* 2004; 44:239–267.
 42. Jaeschke H, Knight TR, Bajt ML. The role of oxidant stress and reactive nitrogen species in acetaminophen hepatotoxicity. *Toxicol. Lett.* 2003; 144:279–288.
 43. Gupta R, Kannan GM, Sharma M and Flora SJS. Therapeutic effects of *Moringa oleifera* on arsenic induced toxicity in rats. *Environ Toxicol Pharmacol.* 2005; 20:456–464.
 44. Geervani P and Devi A. Influence of protein and fat on utilization of carotene from drumstick (*Moringa oleifera*) leaves. *Indian J Med Res.* 1981; 74:548-553.
 45. Siddhuraju P and Becker K. Antioxidant properties of various solvent extracts of total phenolic constituents from three different agroclimatic origins of drumsticks tree (*Moringa oleifera* Lam.) leaves. *J Agric Food Chem.* 2003; 51:44-55.
 46. Wolters-Everhardt E, Dony JM, Lemmens WA, Doesburg WH, De Pont JJ. Buffering capacity of human semen. *Fertil Steril.* 1986; 46:114–9.
 47. Zhou J, Chen L, Li J, Li H, Hong Z, Xie M, et al. The Semen pH Affects Sperm Motility and Capacitation. *PLoS ONE* 2015; 10(7)
 48. Krishna KL, Mruthunjaya K, Patel JA. Antioxidant and hepatoprotective activity of leaf extract of *Justicia gendarussa* burm. *Int. J. Biol. Chem.*, 2009; 3:99-110.