



**CHEMICAL SOCIETY OF NIGERIA
(CSN)**

**34TH
ANNUAL
INTERNATIONAL
CONFERENCE
WORKSHOP &
EXHIBITION**

KWARA 2011

19TH - 23RD SEPTEMBER, 2011

**CONFERENCE
PROCEEDINGS**



**MAIN AUDITORIUM,
UNIVERSITY OF ILORIN,
ILORIN, NIGERIA.**

THEME:
**CHEMISTRY AND
THE MILLENNIUM
DEVELOPMENT
GOALS**

TABLE OF CONTENT (ORGANIC CHEMISTRY PAPERS)

PAGE NO ORG

Eke <i>et al</i>	Methanolysis of Some Agro-Based Triglycerides on the Bed of Cashew Nut Shell	001 – 004
Oyedeji <i>et al</i>	Determination of Anti-Microbial Activity of Extracts of <i>Mitracarpus villosus</i> Based Herbal Dermatitis Soap	005 – 007
Oyedeji <i>et al</i>	Evaluation of Black Soap and Black Soap Admixed with Kaolin for Microbial Activity	008 – 010
Oluwaniyi & Dosumu	Comparative study of the nutrients in <i>Clarias gariepinus</i> (Catfish), <i>Oreochromis niloticus</i> (Tilapia Fish) And <i>Synodontis schall</i> (Kurungu Fish)	011 – 014
Oladoye & Olatunji	Another Labdane Diterpenoid From <i>Aframomum danielli</i> (Zingiberaceae)	015 – 017
Bello & Jimoh	Characterization of Seed and Seed Oil of <i>Syzygium Aromaticum</i>	018 – 010
Owokotomo <i>et al</i>	Comparative Study of the Essential Oils of Seeds and Stem-Bark of <i>Monodora myristica</i> Growing in South Western Nigeria	021 – 023
Arawande & Komolafe	Utilization of Plantain Peel Extract As Antioxidant On Crude Groundnut Oil.	024 – 025
Warra	The Chemistry of Soap Production Utilizing Indigenous Technology and Raw materials	026 – 031
Ibrahim <i>et al</i>	The antimicrobial activities of Crude methanolic extracts of <i>Bauhinia rufescens</i> (Caesalpiniaceae) against some pathogenic microorganisms	032 – 034
Azuaga I.C	Preliminary Phytochemical Screening of Some Selected Medicinal Plants In Chanchanji Community	035 – 036
Obasi <i>et al</i>	Comparative Phytochemical and Antimicrobial Screening of Some Solvent Extracts of <i>Smanaea saman</i> (Fabaceae Or Mimosaceae) Pods	037 – 044
Obasi <i>et al</i>	Antimicrobial and Phytochemical Evaluation of Methanolic Extract and Fractions of <i>Jatropha Curcas</i> Linn (Euphorbiaceae) Stem Bark	045 – 049
Awatefe <i>et al</i>	Characterization of Palm Kernel Shell Powder (Pksp) Using Infrared (Ir) Spectrophotometer	050 – 054
Liman <i>et al</i>	Phytochemical Constituents of the <i>Anacardiaceae</i> [<i>Anacardium occidentale</i> and <i>Mangifera indica</i>]	055 – 057
Odezue Joel	Phytochemical Investigation of the Christmas Cactus <i>epiphyllum</i>	058 – 059

Chinwuba Ike	<i>truncatum</i> For Its Medicinally Active Alkaloids	
Oladosu & Babatunde	Natural Dye and PH Indicator From <i>Sorghum Bicolour</i> Extract	060 – 062
Warra	Saponins As Detergents Additives And Soap Substitute: A Review	063 – 065
Mohammed <i>et al</i>	Proximate, Minerals and Anti-Nutritional Factors of <i>Gardenia aqualla</i> (<i>Gauden Dutse</i>) Fruit Pulp	066 – 070
Yusha'u <i>et al</i>	Phytochemical Screening and Antibacterial Activities of <i>Boswellia dalzielli</i> extracts	071 – 074
Penuel <i>et al</i>	Proximate Composition and Elemental Analysis of <i>Citrullus vulgaris</i> (Guna) Seed	075 – 081
Ghuluze <i>et al</i>	Analysis of Oils Extracted From Two Species of Groundnut Seeds In Yobe State	082 – 084
Yusha'u <i>et al</i>	Phytochemical Screening and Antibacterial Properties of <i>Bridelia ferruginea</i> Extracts	085 – 088
Ukoha <i>et al</i>	Spectrophotometric Determination of Lamivudine, Nevirapine and Zidovudine Using 2, 3-Dichloro-5,6-Dicyano-1,4-Benzoquinone(Ddbq) and Chloranilic Acid	089 – 093
Izuagie <i>et al</i>	Isolation and Characterization of Starch From Christ Thorn Seeds	094 – 098
Danbature <i>et al</i>	The Nutritional Values of Palmyra Tuber Seedling	099 – 100
Gumel <i>et al</i>	Dyeing Properties and Dyeing Kinetics of Colourant Obtained from the Pods of <i>Sienna Siamea</i> L	101 – 104
Fai <i>et al</i>	Extraction and Evaluation of the Quality of Seed Oil of <i>Thevetia neriifolia</i> Just	105 – 107
Daben & Dashak	Extraction, Isolation and Characterization of The Constituents of <i>Crinum Zeylanicum</i>	108 – 110
Gunda <i>et al</i>	Preliminary Phytochemical Analysis and Antimicrobial Activity of <i>Acacia Macrothyrsa</i> Roots	111 – 113
Khan <i>et al</i>	Determination of Mineral Composition, Anti-Microbial Activity and Anti-Nutrient Screening of <i>Cleome gynandra</i> (Cat's Whiskers)	114 – 118
Nwanorh	Active Ingredients In <i>Crinum Ornatum</i>	119 – 121
Ukpong & Akpan,	Phytochemical Screening and Mineral Elements Composition of Cocoyam Inflorescence	122 – 124
Haruna <i>et al</i>	Iridiod and Caffeic Glycoside Ester From the Stem and Leaf of	125 – 129

Stachytarpheta angustifolia (Mill) Vahl Verbenaceae

Oladimeji & Oladosu	Chemical and Biological Assessment of <i>Ceiba Pentadra</i> Spine	130 – 134
Okeke <i>et al</i>	Phytochemical and Elemental Analysis of Leaves of <i>Cassi Alata</i> Plant	135 – 137
Lasisi <i>et al</i>	Phytoconstituents From the Root of <i>Hura crepitans</i> (Euphorbiaceae) and Antibacterial Potency	138 - 143
Okoye	Phytochemical Screening, Nutritional Evaluation and Pharmaceutical Potentials of Seed and Seedcoat of <i>Mucuna pruriens</i> (Velvet Bean)	144 – 148
Ezebuwa & Dashak	Some Properties of A and B Amylase From <i>Eluesine coracana</i> Malt	149 – 154
Ukoha <i>et al</i>	Tannins and Other Phytochemical of the <i>Samanea saman</i> Pods and Their Antimicrobial Activities	155 – 162
Ameen <i>et al</i>	Toxicity and Antimetabolic Properties of Biologically Treated Neem Seed (<i>Azadiracta indica</i>) Kernel Cake	163 – 165
Jigam <i>et al</i>	Toxicological Analysis of the Methanolic Extract of the Stem Bark of <i>Ficus thonningii</i>	166 – 169
Olojede & Ayanbadejo	Phytochemical, Toxicity, Antimicrobial and Antioxidant Screening of Extracts Obtained from <i>Laportea aestuans</i> (Gaud)	170 – 178
Ibrahim <i>et al</i>	The antimicrobial activities of Crude methanolic extracts of <i>Bauhinia rufescens</i> (Caesalpiniaceae) against some pathogenic microorganisms	179 – 181
Kendeson & Dashak	Physico-Chemical, Phytochemical and Mineral Profiles of the Root Bark of <i>Jatropha curcas</i> (Linn) Obtained From Mimyak In Kanke L.G.A of Plateau State.	182 – 184
Orubite - Okorosaye <i>et al</i>	Synthesis Of Trans-3-4-Dibenzyl-2-Oxo-Tetrahydro Furan	185 – 187
Bako <i>et al</i>	Phytochemical screening and antimicrobial characteristics of <i>Senna obtusifolia</i> [<i>Cassia obtusifolia</i> L] seed and leaves	188 – 190
Isah <i>et al</i>	Phytochemical and Antimicrobial Analysis of Stem-Leaf of Methanol Extract of <i>Cochlospermum planchonii</i> Kunth	191 – 194
Adeniyi & Adejoro	Increasing the rate of Novel Drug Discovery through Exploration of Natural Products Biodiversity and Combinatorial Chemistry	195 – 198
Adeniyi & Adejoro	The Structural Based Drug Design Approach and the Roles of Molecular Simulation	199 – 202
Hassan <i>et al</i>	Nutrient composition of a variety of <i>Lagenaria siceraria</i> seed	203 – 206

PRELIMINARY PHYTOCHEMISTRY AND ANTIMICROBIAL PROPERTIES OF *BALERIA OPACA*.

A. A. HAMID^{1*}, O.O. AIYELAAGBE² and A.A. YUSUF¹

¹Department of Chemistry, University of Ilorin, P.M.B. 1515, Ilorin-Nigeria.

E-mail: hamidmemo@yahoo.com; hamid.aa@unilorin.edu.ng; hamidmemo@gmail.com.

²Department of Chemistry, University of Ibadan, Ibadan-Nigeria.

E-mail: oaiyelaagbe@yahoo.com

Corresponding Author's E-mail: hamidmemo@yahoo.com; hamid.aa@unilorin.edu.ng; hamidmemo@gmail.com.

Phone: +2347C35931646

ABSTRACT

The preliminary phytochemical studies of *Baleria opaca* (whole plant) extracts revealed the presence of saponins, tannins, glycosides, alkaloids and anthraquinones, but reducing sugars and flavonoids were not detected in the extracts. The hexane, ethylacetate and methanol extracts effectively inhibited the growth of tested bacteria and fungi at different concentrations respectively. The Methanol extract exhibited higher antibacterial properties than hexane and ethylacetate extracts on *Staphylococcus aureus* and *Bacillus subtilis* (gram positive), *Escherichia coli*, *Pseudomonas aeruginosa*, *Salmonellae typhi* and *Klebsiellae pneumoniae* (gram negative). All the extracts also exhibited significant antifungal properties on *Candida albicans*, *Aspergillus niger*, *Rhizopus stolon*, *penicillium notatum*, *Tricophyton rubrum* and *Epidermophyton floccosum* at concentrations between 25 and 200 mg/ml.

Keywords: *Baleria opaca*, antibacterial, antifungal, Acanthaceae, ethnomedicine.

INTRODUCTION

Medicinal plants have played essential and exceptional roles since early century in the treatment of all kinds of diseases in Africa and other parts of the world. Most Africans still depend on tradition medicine / ethnomedicine of these plants for their medical care because they are considered safer, cheaper, more effective and culturally reliable. Bacterial and fungal infections are prevalent in our society, especially in Africa and these manifests in various diseases like pneumonia, diarrhoea, typhoid, gonorrhoea, syphilis and other sexually transmitted diseases. The search for new, safer and cheaper drugs especially from plants, to cure these diseases is on the increase. *Barleria opaca* (also known as child's vegetable) belong to the family Acanthaceae (Acanthus family) which is a taxon of dicotyledonous flowering plant containing 221 genera and at least 4000 species [1]. Most of them are tropical herbs, shrubs, or twining vines; some are epiphytes. Only a few species are distributed in temperate regions. Their main centres of distribution are Indonesia and Malaysia, Africa, Brazil and Central America. The representatives of the family can be found in nearly every habitat, including dense or open forests, in scrublands, on wet fields and valleys, at the sea coast and in marine areas, and in swamps and as an element of mangrove woods [2]. *B. opaca* is a scrambling shrub with hairy stems and sometimes rooting at the nodes. leaves opposite, simple; petiole up to 0.5cm long, blade elliptical, 7.5 by 3.5cm, tapering at both ends with scattered simple hairs at both surfaces. The plant occurs in Western and Central tropical Africa, from Cote d'Ivoire to Gabon. The leaves of *B. opaca* are used to treat children for piles by squatting in a warm decoction. In Nigeria the whole plant is used in treating jaundice,

rheumatism and paralysis, and the leaf sap is applied against catarrh [3]. The pharmacological and phytochemistry of *B. opaca* have not been reported. It is on the basis of these, that we investigate the phytochemicals and antimicrobial properties of the plant.

MATERIALS AND METHODS

Collection and authentication of the plant material.

The plant material of *Baleria opaca* was collected from Ibadan, Oyo State of Nigeria, October 2010. Botanical identification and authentication was done by Mr. A.W. Ekundayo of the Forestry Research Institute of Nigeria (FRIN), Ibadan, Nigeria where a voucher specimen (FHI108470) was deposited.

Preparation of plant extracts

The whole plant of *Baleria opaca* was air-dried, grounded and weighed (689g). The dried plant was successively extracted in hexane, ethylacetate and methanol for 10 days respectively using cold extraction method. Hexane (5g), ethylacetate (5.5g) and methanol (7g) extracts were obtained by evaporation and stored in the refrigerator for further use.

Phytochemical studies

The Preliminary phytochemical screening of the hexane, ethylacetate and methanol extracts of *B. coccineus* leaves was done using standard procedures [4, 5, 6, 7, and 8].

1) Saponins: Small quantity of each extract was boiled with 5 ml of distilled water, filtered and cooled.

a). Frothing: To the filtrate (2.5 ml) about 10 ml of distilled water was added and shaken vigorously for 2 minutes. Frothing observed indicates a positive test.

b). Emulsification: To the filtrate (2.5 ml) added 3 drops of olive oil and shaken vigorously for 2 minutes. An emulsified layer indicates a positive test.

2) **Alkaloids:** Small quantity of each extract was stirred with 5 ml of 1% hydrochloric acid for five minutes on a water bath and then filtered. Of the filtrate of each extract was divided into two portions. Mayer's reagent was added to one portion; occurrence of creamy white precipitate was taken as positive. To the second portion few drops of Dragendorff's reagent was added and appearance of orange red precipitate was regarded as positive for the presence of alkaloids.

3) **Glycosides (Keller-killiani Test):** Small quantity of each extract was diluted in 5 ml of distilled water. Added 2 ml of glacial acetic acid containing one drop of ferric chloride solution (3.5%) to each. This was underlay with 1 ml of concentrated sulfuric acid. A radish brown ring is formed at the interface and upper layer turns bluish green on standing indicates the presence of a deoxy sugar characteristic of cardiac glycosides.

b) **Method-2:** Small quantity of each extract moistened with 5 ml in distilled water and filtered.

Few drops of chloroform were added to each (to enhance enzymatic activity). A sodium picrate-saturated filter paper strip was hanged at the neck of the flask with the help of the cork and warmed the flask. The filter paper strip turned brick-red or maroon is indicated the presence of cyanogenetic glycosides.

4) **Tannins:**

a) **Ferric Chloride Test:** Small quantity of each extract was boiled in 10 ml of water in a test tube and then filtered while hot and a few drops of 0.1% ferric chloride solution were added to the filtrate. A brownish green or a blue-black coloration indicates as a positive test.

b) **Lead Acetate Test:** Small quantity of each extract was taken in a test tube and diluted with 5 ml of distilled water. Add few drops of a 1% solution of lead acetate to each. A yellow or red precipitate indicates a positive test.

5) **Flavonoids:** Three methods were used to determine the presence of flavonoids in the extracts.

a) **Method-1:** Dilute ammonia solution (5 ml) was added to aqueous filtrate of each extract followed by addition of concentrated H₂SO₄ acid (1 ml). A yellow colouration that disappears on standing indicates the presence of flavonoids.

b) **Method-2:** Few drops of 1% aluminium solution were added to aqueous filtrate of the each extract. A yellow coloration indicates the presence of flavonoids.

c) **Method -3:** A small portion of the each extract was heated with 10 ml of ethyl acetate over a steam bath for 3 min. The mixture was filtered and 4 ml of the filtrate was shaken with 1 ml of dilute ammonia solution. A yellow coloration indicates the presence of flavonoids.

6) **Steroids:**

a) **Liebermann-burchard's Test:** Small amount of each extract was dissolved in 1 ml of chloroform. Added 2 ml of acetic anhydride and 1 ml of concentrated H₂SO₄ acid to each portion. A greenish color is produced which turns blue on standing indicates the presence of steroids.

b) **Salkowski's Test:** Small amount of the extract was dissolved in 2ml of chloroform. Concentrated sulphuric acid was carefully added to a lower layer. A reddish-brown colour at the interphase indicates the presence of deoxysugar characteristics of cardenolides. A violet ring may form just above the ring and gradually spread throughout the layer.

7) **Reducing sugars (Fehling's Test):** A small portion of each of the extract was shaken with distilled water and filtered. The filtrate was boiled with drops of Fehling's solution A and B for 2 minutes. An orange-red precipitate on boiling with Fehling's solution indicates the presence of reducing sugars.

8) **Anthraquinones:** A small portion of each extract was boiled with 10 ml of sulfuric acid, traces of ferric chloride solution was added and filtered while hot. The filtrate was shaken with 5 ml of chloroform. The chloroform layer was taken into another test tube and 1 ml of dilute ammonia was added to each portion. Rose-pink color in the aqueous layer indicates the presence of anthraquinones.

Antimicrobial Assay

Microorganisms: Cultures of six human pathogenic bacteria made up of four gram negative and two gram positive were used for the antibacterial assay. These were; *Salmonella typhii* (UCH 4801), *Escherica coli* (UCH 0026), *Pseudomonas aeruginosa* (UCH 1102) and *Klebsiellae pneumoniae* (UCH 2894) belongs to the gram-negative and *Bacillus subtilis* (UCH 7423) while *Staphylococcus aureus* (UCH 2473) belongs to the gram-positive. For the Antifungal assay, six fungi were also utilized. These were; *Candida albicans*, *Aspergillus niger*, *Rhizopus stolon*, *Penicillum notatum*, *Tricophyton rubrum* and *Epidermophyton floccosum*. All the microorganisms used were clinical strains from the Medical Microbiology (University College Hospital, Ibadan) and screened in the Laboratory of Pharmaceutical Microbiology Department, University of Ibadan, Ibadan, Nigeria.

Media: Nutrient agar, Sabouraud dextrose agar, nutrient broth and tryptone soya agar were used in this study. Hexane, ethylacetate and methanol were also used in solubilizing the extracts and as negative controls in the assays.

Antimicrobial agents: Gentamycin (10 µg/mL) and Tioconazole (0.7 mg/mL) were included as standard reference drugs in the study.

Antimicrobial activity determination

Agar diffusion-pour plate method (bacteria): An overnight culture of each organism was prepared by taken two wireloop of the organism from the stock and inoculated each into the sterile nutrient broth of 5ml, each incubated for 18-24hr at 37°C. From overnight culture, 0.1 mL of each organism was taken and put into the 9.9mL of sterile distilled water to obtained 10⁻² inoculum concentration of the organism.

From the diluted organism (10⁻²), 0.2mL was taken into the prepared sterile nutrient agar cooled to about 40-45°C, then poured into sterile Petri dishes.

to solidify for about 45-60min. Using a sterile cork-borer of 8mm diameter, the wells were made according to the number of the test tubes for the experiment. For this work 8 wells were made. The graded concentrations of the extracts were put into the wells accordingly including the controls. The studies were done in duplicates to ascertain the results obtained. The plates were left on the bench for about 2hrs to allow the extract diffuse properly into the nutrient agar i.e. pre-diffusion. The plates were incubated for 18-24hr at 37°C.

Agar diffusion-surface plate method (fungi): A sterile sabouraud dextrose agar was prepared accordingly and aseptically poured into the sterile plates in triplicates and solidified properly. 0.2mL of the 10^{-2} inoculum concentration of the organism was spread on the surface of the agar using a sterile Petri-dish cover to cover all the surface of the agar. Eight wells were bored using a sterile

cork-borer of 8mm diameter. The graded concentrations of the extracts were put into the including the controls. All the plates were left on the bench for 2hr to allow the extract diffuse properly into the agar i.e. pre-diffusion. The plates were incubated at 25°C for 72hr [9,10,11].

RESULTS AND DISCUSSIONS

The preliminary phytochemical studies of the hexane, ethylacetate and methanol extracts of *Baleria opaca* revealed the presence of tannins, glycosides and anthraquinones (Table 1).

Saponins and alkaloids were detected in ethylacetate and methanol extracts of the plant but the two metabolites were absent in the hexane extract. The presence of steroids was established only in methanol extract. However, reducing sugars and flavonoids were not found in all the extracts.

Table 1: Phytochemical constituents of the hexane, ethylacetate and methanol extracts of *Baleria opaca* (whole plant).

Secondary metabolites	Extracts (whole plant)		
	Hexane	Ethylacetate	Methanol
Alkaloids	-	++	++
Saponins	-	++	++
Tannins	++	++	++
Reducing sugars	-	-	-
Steroids	-	-	++
Glycosides	++	++	++
Flavonoids	-	-	-
Anthraquinones	++	++	++

Key: - Absent ++ Present

Anti bacteria activities of hexane, ethylacetate and methanol extracts is presented in Table 2. In the table, methanol extract showed higher inhibition on the six tested organisms than the Hexane and ethylacetate extracts. However, all the bacteria strains were sensitive to the three extracts at concentrations ranging from 12.5 to 200 mg/mL. Hexane and methanol extracts exhibited higher inhibition against the growth of *Staphylococcus aureus*, *Escherichia coli*, *Bacillus subtilis*, *Pseudomonas aeruginosa* and *Klebsiellae pneumoniae* than ethylacetate extract of the plant. Meanwhile, the methanol extract showed higher antibacterial properties on the six tested organisms than both the hexane and ethylacetate extracts.

The result of the antifungal activities of the hexane, ethylacetate and methanol extracts of *B. opaca* at concentrations between 6.25 and 200mg/mL is presented in Table 3.

In the table, *Candida albicans*, *Aspergillus niger*, *Rhizopus stolon*, *Penicillium notatum*, *Tricophyton rubrum* and *epidermophyton floccosum*. *Candida albicans* showed higher sensitivity to hexane and ethylacetate extracts than methanol extract, while methanol extract had higher antifungal activities on *Aspergillus niger* and *Tricophyton rubrum* than both hexane and ethylacetate extracts of the plant. Further more, ethylacetate showed higher inhibition on *Rhizopus stolon* and *Epidermophyton floccosum* than *Aspergillus niger*, *Penicillium notatum* and *Tricophyton rubrum* while hexane extract showed higher inhibition on *Candida albicans* and *Aspergillus niger* than other test fungi at concentrations between 25 and 200mg/mL. However, the sensitivity of the test bacteria and fungi to all the extracts were concentration dependent, activity being higher at higher concentration of the extracts.

Table 2: Antibacterial activities of the hexane, ethylacetate and methanol extracts of *Baleria opaca* (whole plant).

Extracts	Extract conc/Ref./ Control (mg/ml)	Diameter of well = 8 mm Diameter of zone of inhibition of bacteria (mm)					
		S.a	E.coli	B.sub	Ps.a	Kleb	Sal.
Hexane	6.25	—	—	—	—	—	—
	12.5	—	—	—	10	—	—
	25	10	—	10	14	10	—
	50	12	10	12	16	12	10
	100	14	14	14	18	14	12
	200	16	18	18	20	16	14
	Hexane	—	—	—	—	—	—
Ethylacetate	Gentamycin	38	36	34	34	36	38
	6.25	—	—	—	—	—	—
	12.5	—	—	—	—	—	—
	25	—	—	10	—	—	10
	50	10	10	12	10	12	12
	100	12	12	14	12	14	14
	200	14	14	16	14	16	18
Methanol	Ethylacetate	—	—	—	—	—	—
	Gentamycin	38	36	36	34	36	34
	6.25	—	—	—	10	—	—
	12.5	10	10	—	14	10	10
	25	12	12	10	16	12	12
	50	14	14	14	18	14	14
	100	16	18	16	20	16	18
	200	20	22	20	24	18	20
	Methanol	—	—	—	—	—	—
	Gentamycin	37	33	36	34	32	34

Key: S.a *Staphylococcus aureus* C.a *Candida albicans*
E.coli *Escherichia coli* A.n *Aspergillus niger*
B.sub *Bacillus subtilis* Rhiz *Rhizopus stolon*
Ps.a *Pseudomonas aeruginosa* Pen *Penicillium notatum*
Kleb *Klebsiella pneumoniae* T.r *Tricophyton rubrum*
Sal *Salmonella typhi* E.f *Epidermophyton floccosum*

Table 3: Antifungal activities of the hexane, ethylacetate and methanol extracts of *Brysocarpus coccineus* Leaves.

Extracts	Extract conc/Ref./ Control (mg/ml)	Diameter of well = 8 mm Diameter of zone of inhibition of bacteria (mm)					
		C.a	A.n	Rhiz	Pen	T.r	E.f
Hexane	6.25	—	—	—	—	—	—
	12.5	—	—	—	—	—	—
	25	10	—	—	—	—	—
	50	12	10	10	—	—	10
	100	14	12	12	10	10	12
	200	16	16	14	12	12	14
	Hexane	—	—	—	—	—	—
Ethylacetate	Tioconazole	26	24	20	24	26	26
	6.25	—	—	—	—	—	—
	12.5	—	—	—	—	—	—
	25	10	—	10	—	—	10
	50	12	—	12	10	—	12
	100	14	10	14	12	10	14
	200	18	12	16	14	12	16
Methanol	Ethylacetate	—	—	—	—	—	—
	Tioconazole	26	26	24	24	24	26
	6.25	—	—	—	—	—	—
	12.5	—	—	—	—	—	—

	25	—	10	—	10	—
	50	—	12	—	10	12
	100	10	14	10	12	14
	200	12	16	12	14	18
	Methanol	—	—	—	—	—
	Tioconazole	26	24	25	22	22
Key:	S.a	<i>Staphylococcus aureus</i>	E.coli	<i>Escherichia coli</i>		
	B.sub	<i>Bacillus subtilis</i>	Ps.a	<i>Pseudomonas aeruginosa</i>		
	Kleb	<i>Klebsiellae pneumoniae</i>	Sal	<i>Salmonellae typhi</i>		
	C.a	<i>Candida albicans</i>	A.n	<i>Aspergillus niger</i>		
	Rhiz	<i>Rhizopus stolon</i>	Pen	<i>Penicillium notatum</i>		
	T.r	<i>Tricophyton rubrum</i>	E.f.	<i>Epidermophyton floccosum</i>		

CONCLUSIONS

The fungicidal and bactericidal properties of *Baleria opaca* extracts suggest the use of the plant for the treatment of infectious diseases caused by the tested organisms. Isolation and characterization of bioactive compounds that are responsible for the observed biological activities is ongoing.

ACKNOWLEDGEMENTS

H.A.A is grateful to the Department of Chemistry, University of Ilorin, Nigeria for the vital assistance rendered during this research.

REFERENCES

1. Bentham, G. and Hooker, J. D. (1876). *Genera Plantarum* 2: 1060-1122. London: Reeve.
2. Daniel, T. F. (1983). Systematics of Holographis (Acanthaceae). *Journal of the Arnold Arboretum* 64: 129-160.
3. Irvine, F.R. (1961). *Woody plants of Ghana, with special references to their uses*. Oxford University Press, London. pp 868.
4. Edeoga, H.O., Okwu, D.E., Mbaebie, B.O. (2005). Phytochemical constituents of some Nigerian medicinal plants. *Afr. J. Biotech.*, 4, 685-688.
5. Harborne, J.P. (1991). *Phytochemical Methods*: 2nd ed. Oxford University Press, London, 51.
6. Harborne, J.P. (1998). *Phytochemical Methods: A guide to modern technique of plant analysis*, 2nd ed. Chapman and Hall, London.
7. Trease, G.E., Evans, W.C. (1983). *Drugs of Biological Origin*. In *Pharmacognosy* 12th ed. United Kingdom: Balliere Tindall. 309-540.
8. Trease, G.E., Evans, W.C. (1989). *Pharmacognosy*. Brailiar Tiridel can, 13th ed. Macmillian Publishers.
9. Van den Berge, A.D., Vlietinck, A.J. (1991). *Methods in plants biochemistry*. Vol. 6. Assays for bioactivity. Hostettmann K. Ed. Academic press, London. 47-69.
10. Kavanagh, F. (1972). *Analytical microbiology*, Academic Press, New York
11. Zwadyk, P. (1972). *Enteriobactericeae*, in *Zieser microbiology*, 20th Ed. Pp 544 – 545, Georg Thieme verlag, Stuttgart.